

Using Redundant Parameterizations to Fit Hierarchical Models

Andrew GELMAN, David A. VAN DYK,
Zaiying HUANG, and W. John BOSCARDIN

Hierarchical linear and generalized linear models can be fit using Gibbs samplers and Metropolis algorithms; these models, however, often have many parameters, and convergence of the seemingly most natural Gibbs and Metropolis algorithms can sometimes be slow. We examine solutions that involve reparameterization and overparameterization. We begin with parameter expansion using working parameters, a strategy developed for the EM algorithm. This strategy can lead to algorithms that are much less susceptible to becoming stuck near zero values of the variance parameters than are more standard algorithms. Second, we consider a simple rotation of the regression coefficients based on an estimate of their posterior covariance matrix. This leads to a Gibbs algorithm based on updating the transformed parameters one at a time or a Metropolis algorithm with vector jumps; either of these algorithms can perform much better (in terms of total CPU time) than the two standard algorithms: one-at-a-time updating of untransformed parameters or vector updating using a linear regression at each step. We present an innovative evaluation of the algorithms in terms of how quickly they can get away from remote areas of parameter space, along with some more standard evaluation of computation and convergence speeds. We illustrate our methods with examples from our applied work. Our ultimate goal is to develop a fast and reliable method for fitting a hierarchical linear model as easily as one can now fit a nonhierarchical model, and to increase understanding of Gibbs samplers for hierarchical models in general.

Key Words: Bayesian computation; Blessing of dimensionality; Markov chain Monte Carlo; Multilevel modeling; Mixed effects models; PX-EM algorithm; Random effects regression; Redundant parameterization; Working parameters.

Andrew Gelman is Professor, Department of Statistics and Department of Political Science, Columbia University, New York (E-mail: gelman@stat.columbia.edu; <http://www.stat.columbia.edu/~gelman/>). David van Dyk is Professor, Department of Statistics, University of California, Irvine, (E-mail: dvd@ics.uci.edu; <http://www.ics.uci.edu/~dvd/>). Zaiying Huang is Statistician, Circulation Department, *New York Times* (E-mail: huangz@nytimes.com). W. John Boscardin is Associate Professor, Department of Biostatistics, University of California, Los Angeles (E-mail: jbosco@ucla.edu).

© 2008 *American Statistical Association, Institute of Mathematical Statistics, and Interface Foundation of North America*

Journal of Computational and Graphical Statistics, Volume 17, Number 1, Pages 95–122
DOI: 10.1198/106186008X287337

1. INTRODUCTION

1.1 BACKGROUND

Hierarchical linear models (also called multilevel models, random effects regressions, and mixed effects models) are important tools in many areas of statistics (for reviews see, e.g., Robinson 1991; Longford 1993; Goldstein 1995). In recent years, there has been increasing interest in the Bayesian formulation (Lindley and Smith 1972), which accounts for the uncertainty in the variance parameters, a feature that is particularly important when the hierarchical variances are hard to estimate or to distinguish from zero (see Carlin and Louis 2000 and Gelman et al. 1995 for discussions and examples). Modern Bayesian inference generally entails simulating draws of the parameters from their posterior distribution. For hierarchical linear models, this can be done fairly easily using the Gibbs sampler (see Gelfand and Smith 1990). For hierarchical generalized linear models, essentially the same algorithm can be used by linearizing the likelihood and then applying a Metropolis-Hastings accept/reject rule at each step (see Gelman et al. 1995; Gilks, Richardson, and Spiegelhalter 1996).

Unfortunately, the standard algorithms for computing posterior simulations from hierarchical models can have convergence problems, especially when hierarchical variance components are near zero. Gibbs samplers (and also EM, Metropolis, and other algorithms) can get stuck because of dependence in the posterior distribution between batches of coefficients and their variance parameters. In this article, we consider computational methods that aim to improve convergence by embedding the target posterior distribution into a larger model. *Auxiliary variables* (Besag and Green 1993) and *data augmentation* (Tanner and Wong 1987) are two common methods that work in this way. Auxiliary variables are added to a model in such a way that conditioning on these variables allows a number of model parameters that would otherwise be updated one at a time to be blocked into one joint draw in a Gibbs sampler. Data augmentation, on the other hand, introduces additional variables (called augmented data or latent variables) to simplify the conditional draws of the Gibbs sampler. For example, the complete conditional distributions may become standard distributions in the larger model that includes the augmented data. Together these methods can simplify and reduce dependence in the conditional specifications in a joint posterior distribution.

In this article, we focus on a third strategy that builds on data augmentation. *Parameter expansion* introduces a specific sort of variable known as a working parameter. Such parameters are identifiable given the observed data and augmented data, but are not identifiable given the observed data alone. This partial identifiability allows additional flexibility in constructing samplers. With care, it can be shown that this flexibility can be used to improve the convergence properties of the samplers. In this way, we are able to add structure that can be leveraged for improved computation without altering the fitted observed data model. The method was introduced by Meng and van Dyk (1997) and Liu, Rubin, and Wu (1998) for the EM algorithm and extended to the data augmentation algorithm by Meng and van Dyk (1999) and Liu and Wu (1999). Meng and van Dyk (1999) introduced the terms *conditional* and *marginal* augmentation to indicate whether the working parameter

is conditioned on or averaged over in the iteration; conditional augmentation essentially indexes a family of transformations by the working parameters. Here we compare and extend the new algorithms to hierarchical linear and generalized linear models (see also van Dyk 2000; van Dyk and Meng 2001; Imai and van Dyk 2004). We evaluate our various proposed algorithms from the standpoint of computation time per iteration, number of iterations required for approximate convergence, and the autocorrelation in the resulting chains.

This article proceeds as follows. The rest of Section 1 contains background material including discussion of the basic hierarchical linear models, standard Gibbs/Metropolis algorithms based on scalar or vector updating using linear regression computations, structured Markov chain Monte Carlo, and conditional and marginal augmentation. Section 2 presents two improvements in computation: the parameter-expanded hierarchical model and the scalar updating algorithm with rotation, an example of conditional augmentation. Section 3 presents theoretical arguments on the computation time per iteration and convergence times for several proposed Gibbs sampler implementations, including an innovative argument based on how quickly a Markov chain can free itself from getting stuck in remote areas of the parameter space. Section 4 presents two examples, Section 5 discusses extensions to generalized linear model, and Section 6 concludes with recommendations and ideas for further research.

1.2 NOTATION FOR HIERARCHICAL LINEAR MODELS

1.2.1 Hierarchical Linear Model

In setting up any mathematical notation, there is a tension between conciseness and generality. To keep the focus in this article on the basic computational ideas, we work with a relatively simple form of the Gaussian hierarchical linear regression model with a vector of data y and a vector of coefficients β that are arranged in batches:

$$y|\beta \sim N(X\beta, \Sigma_y) \quad (1.1)$$

$$\beta \sim N(\beta_0, \Sigma_\beta). \quad (1.2)$$

Hierarchical linear models with more than two levels can be expressed in this canonical form by extending the vector β to include the higher-level linear parameters in the hierarchy and augmenting y and X appropriately (see Gelman et al. 1995; Hodges 1998; Sargent, Hodges, and Carlin 2000). For simplicity, we assume the mean vector β_0 is known.

The vector β can include modeled and unmodeled parameters (sometimes called “fixed and random effects,” terminology we avoid for reasons described by Gelman (2005, sec. 6)). Elements of β that are unmodeled (e.g., in a multilevel model, individual-level coefficients that do not vary by group) can be combined into a batch whose group-level variance is infinite. The notation could equivalently be developed with modeled and unmodeled parameters treated separately (as in Laird and Ware 1982).

A key aspect of the model is the parameterization of the variance matrices, Σ_y and Σ_β . We shall assume that each Σ_z is diagonal with a vector of unknown elements $\sigma_z^2 = (\sigma_{zk}^2 :$

$k = 1, \dots, K_z$), where we are using z as a floating subscript to stand for y or β :

$$\begin{aligned}\Sigma_y &= \text{Diag}(W_y \sigma_y^2) \\ \Sigma_\beta &= \text{Diag}(W_\beta \sigma_\beta^2),\end{aligned}\tag{1.3}$$

and W_y and W_β are indicator matrices of 0's and 1's, with exactly one 1 in each row. W_y and W_β are not, in general, square: they indicate the batching of the data and parameters. Each component of σ_y and σ_β is the standard deviation assigned to a subset, or batch, of the components of y or β . The data are partitioned into K_y batches, the regression coefficients are partitioned into K_β batches, and the columns of W_y and W_β correspond to vectors of indicator variables for each of the batches. (Depending on the details of the model, the actual computation may be done by selecting appropriate batches of coefficients rather than literally by matrix multiplication. The matrix notation, however, has the advantage of automatically including nonnested groupings which could not simply be handled by grouping the elements of β into ordered batches.)

1.2.2 Prior Distributions on the Variance Parameters

We assign inverse- χ^2 prior distributions to the variance parameters:

$$\begin{aligned}\sigma_{yk}^2 &\sim \text{Inv-}\chi^2(v_{yk}, \sigma_{0yk}^2), \quad \text{for } k = 1, \dots, K_y \\ \sigma_{\beta k}^2 &\sim \text{Inv-}\chi^2(v_{\beta k}, \sigma_{0\beta k}^2), \quad \text{for } k = 1, \dots, K_\beta.\end{aligned}\tag{1.4}$$

The prior degrees of freedom ν and scale parameters σ_0 can either be preset or assigned a noninformative prior distribution. There are several important special cases of this choice of prior distribution. First, the standard noninformative prior distribution, $p(\sigma_y, \sigma_\beta) \propto \prod_{k=1}^{K_y} \sigma_y^{-1}$, corresponds to $(v_{yk} = 0, \sigma_{0yk} = 0)$ for $k = 1, \dots, K_y$, and $(v_{\beta k} = -1, \sigma_{0\beta k} = 0)$ for $k = 1, \dots, K_\beta$. Second, variance components with known values correspond to $\nu_{zk} = \infty$ and are, of course, not altered in any of the Gibbs sampler algorithms. Third, components of β with noninformative flat prior distributions are associated with variance components $\sigma_{\beta k}$ that are fixed at ∞ . Fourth, components of β with fixed or known values are associated with variance components $\sigma_{\beta k} = 0$. See Gelman (2006) for further discussion of prior distributions for hierarchical models.

1.2.3 Extensions of the Models

The model can be extended in various ways, including nondiagonal variances (Gelman et al. 1995), multivariate clustering of regression parameters (Barnard, McCulloch, and Meng 1996; Daniels and Kass 1999), informative prior distributions on the regression coefficients, unequal variances (Boscardin and Gelman 1996), robust models, and many other variants in the statistics and econometrics literature, perhaps most notably models for variable selection and model averaging (Raftery 1996). Most of these extensions can be fit into our framework by adding additional parameters and thus additional steps in the Gibbs or Metropolis algorithms. Rather than trying to present an ideal algorithm to handle all contingencies, our goal is to understand what works with an important and nontrivial

basic family of models. In Section 5 we discuss extensions of our methods and algorithms to a class of generalized linear models.

1.3 BASIC GIBBS SAMPLERS FOR THE HIERARCHICAL NORMAL MODEL

There are two simple Gibbs sampler algorithms for the hierarchical linear model. In both algorithms, the variance parameters $\sigma = (\sigma_y, \sigma_\beta)$ are drawn from their joint distribution given β and y . The algorithms differ in their sampling of β : in one version the components of β are drawn one at a time, and in the other version the vector β is drawn all at once. As Sargent, Hodges, and Carlin (2000) showed, aspects of these two algorithms can be combined for greater efficiency; we discuss this further in Section 2. When we speak of an *iteration* of either algorithm, we refer to a single update of all of the free parameters in the model.

The *all-at-once* algorithm draws the vector β from $p(\beta|\sigma_y, \sigma_\beta, y)$ by running the regression of y_* on X_* with variance matrix Σ_* , where

$$y_* = \begin{pmatrix} y \\ \beta_0 \end{pmatrix}, \quad X_* = \begin{pmatrix} X \\ I \end{pmatrix}, \quad \text{and} \quad \Sigma_* = \begin{pmatrix} \Sigma_y & 0 \\ 0 & \Sigma_\beta \end{pmatrix} \quad (1.5)$$

combine the data and prior information (Gelman et al. 1995; see also Dempster, Rubin, and Tsutakawa 1991 and Hodges 1998). Given Σ_* , the regression computation yields an estimate of the regression coefficient and an estimate of its variance matrix, which we label $\hat{\beta}$ and V_β , with the understanding that both are functions of the data and the variance parameters. To simulate β in the Gibbs sampler we need not compute V_β explicitly; rather, we can compute the QR decomposition $\Sigma_*^{-1/2} X_* = QR$ and then draw

$$\text{all-at-once Gibbs update:} \quad \beta = \hat{\beta} + R^{-1}z, \quad (1.6)$$

where z is a vector of independent standard normal random variables. (We sometimes write $\hat{\beta}(\sigma)$ and $R(\sigma)$ to emphasize the dependence of $\hat{\beta}$ and R on the variance components.)

The *one-at-time* algorithm samples β componentwise, conditional on all the other parameters:

$$\text{one-at-a-time Gibbs update: for each } j, \text{ sample } \beta_j \sim p(\beta_j|\beta_{-j}, \Sigma_y, \Sigma_\beta, y), \quad (1.7)$$

where β_{-j} is all of β except β_j . The conditional distribution in expression (1.7) is a simple univariate normal distribution.

The one-at-a-time computations are slightly more difficult to set up in the general case than the vector Gibbs sampler, but they have the advantage of never requiring large matrix operations. If the updating step (1.7) is set up carefully, with the appropriate intermediate results held in storage, this algorithm can be very efficient in terms of computation time per iteration (Boscardin 1996). There is also the potential for further speeding the scalar computations by taking advantage of zeroes in the X matrix.

For either algorithm, updating the variance components is simple; their conditional posterior distributions, given β , are independent,

$$\sigma_{zk}^2|\beta \sim \text{Inv-}\chi^2\left(v_{zk} + n_{zk}, \frac{v_{zk}\sigma_{0zk}^2 + SS_{zk}}{v_{zk} + n_{zk}}\right),$$

where n_{zk} is the number of components of z that correspond to the variance parameter σ_{zk} (i.e., the sum of the elements of the k th column of W_z), and SS_{zk} is the k th component of the appropriate vector of residual sum of squares:

$$SS_y = W_y \text{Diag}((y - X\beta)\Sigma_y^{-1}(y - X\beta)^t),$$

or

$$SS_\beta = W_\beta \text{Diag}((\beta - \beta_0)\Sigma_\beta^{-1}(\beta - \beta_0)^t), \quad (1.8)$$

with the variance matrices Σ_y and Σ_β defined in (1.3).

1.4 STRUCTURED MARKOV CHAIN MONTE CARLO

If the components of β are highly correlated in their posterior distribution, then the one-at-a-time Gibbs sampler can move slowly. The vector Gibbs sampler avoids this problem by sampling all the components of β at once, but this algorithm can be slow in computation time because it requires a full linear regression computation at each step. (We formalize this run-time computation in Section 3.1.)

This tradeoff is well known and was discussed at length by Sargent, Hodges, and Carlin (2006), who advocated the all-at-once Gibbs sampler to avoid a slowly converging chain. To reduce the per-iteration computational costs, they suggest replacing Σ_* by an approximate value when updating β . The approximation may be updated periodically during the pre-convergence burn-in period, but at some point it is fixed and no longer updated to preserve the Markov property of the chain. The approximation to Σ_* is used in the jumping rule of a Metropolis–Hastings sampler, thus preserving the target posterior distribution as the stationary distribution of the sampler.

Optimally, we would combine the computational speed of the one-at-a-time algorithm with the fast convergence of the all-at-once algorithm. We pursue two strategies that aim to accomplish this without a Metropolis–Hastings correction, at least in the hierarchical linear model. Our strategies are based on the methods of conditional and marginal augmentation, and we conclude this section with a general introduction and example of these methods.

1.5 CONDITIONAL AND MARGINAL AUGMENTATION

Conditional and marginal augmentation (Meng and van Dyk 1999) are a set of techniques that alter the joint distribution of the observed data and the augmented data in order to improve computation. This is done while preserving the marginal distribution of the observed data and thus the target posterior distribution. In the hierarchical linear model, β is treated as augmented data, and we consider alternative forms of $p(y, \beta|\sigma)$ that preserve $p(y|\sigma)$. We alter the joint distribution in a very particular way, and it can be shown that the resulting joint posterior distribution has the target posterior distribution of the parameter as its marginal distribution but has less posterior correlation between the parameter and the augmented data (Meng and van Dyk 1999; van Dyk and Meng 2001). We can take advantage of this reduced correlation to construct a Markov chain with reduced autocorrelation, but with the target posterior distribution of the model parameters as its stationary distribution.

In order to alter the joint distribution of y and β , we introduce a one-to-one transformation $\zeta = \mathcal{D}_\alpha(\beta)$ that is parameterized in terms of the so-called working parameter α . A working parameter indexes a class of joint distributions of the observed and augmented data given the model parameters in such a way that the marginal distribution of the observed data does not depend on the working parameter. In particular, the working parameter indexes a class of equivalent models, in the sense that

$$\int p(y, \zeta | \sigma, \alpha) d\zeta = p(y | \sigma) \quad \text{for all } \alpha \in \mathcal{A};$$

that is, $p(y | \sigma, \alpha)$ does not depend on α . The working parameter does not affect inference based on $p(y | \sigma)$. We use the notation ζ to emphasize that we are using a different joint distribution of the observed and augmented data, with the understanding that this does not affect the marginal distribution of the observed data. In the context of the hierarchical linear model we transform from β to ζ via premultiplication by a matrix depending on α that rescales and potentially decorrelates the components of β .

Conditional and marginal augmentation differ in their treatment of working parameters. Conditional augmentation constructs a Gibbs sampler based on the conditional distributions of

$$p(\zeta, \sigma | y, \alpha) \propto p(y | \zeta, \sigma, \alpha) p(\zeta | \sigma, \alpha) p(\sigma),$$

where, as usual, we assume the working and model parameters are a priori independent. The working parameter is chosen to optimize or at least to improve convergence properties. It is this step that effectively reduces the posterior correlation between the parameters and the augmented data. In contrast, marginal augmentation averages over the working prior density, $p(\alpha)$, and uses a Gibbs sampler based on the conditional distributions of

$$\int p(\zeta, \sigma, \alpha | y) d\alpha \propto p(\sigma) \int p(y | \zeta, \sigma, \alpha) p(\zeta | \sigma, \alpha) p(\alpha) d\alpha.$$

By eliminating the conditioning on α that is implicit in a standard Gibbs sampler, we increase the conditional posterior variance of the augmented data. This in turn allows bigger jumps and reduces the autocorrelation of the Markov chain. We illustrate how to set up and use conditional and marginal augmentation samplers in our hierarchical models in Section 2, illustrating with a simple example in Section 1.6.

When using marginal augmentation, the choice of the working prior density, $p(\alpha)$, affects the convergence behavior of the algorithm; generally more diffuse densities improve mixing. Unfortunately, improper working prior densities lead to nonpositive recurrent Markov chains since $p(\alpha | y) = p(\alpha)$ and hence $p(\zeta, \sigma, \alpha | y)$ is improper. With judicious choice of the improper working prior density, however, we can ensure that the subchain corresponding to the model parameters is recurrent with stationary distribution $p(\sigma | y)$; see Meng and van Dyk (1999) and Liu and Wu (1999). The basic strategy is to construct a sequence of Markov transition kernels on σ or (β, σ) each with the target posterior distribution as their stationary distribution. The sequence is further constructed so that its limiting transition kernel is the proper Markovian kernel constructed with the improper working prior density. This strategy is illustrated in the Appendix.

1.6 EXAMPLE: A SIMPLE HIERARCHICAL MODEL

We illustrate the basic idea in a simple example to which we shall return in Section 4.1: a hierarchical linear model of the form,

$$y_j | \mu, \beta_j \stackrel{\text{indep.}}{\sim} N(\mu + \beta_j, \sigma_j^2), \quad \text{with } \beta_j \stackrel{\text{iid.}}{\sim} N(0, \sigma_\beta^2), \quad (1.9)$$

with σ_j known for each j . A Gibbs sampler for this model can get stuck near $\sigma_\beta = 0$: once this group-level variance is set near zero in the chain, the β_j 's are likely to be drawn near zero in their step of the Gibbs updating, at which point σ_β is likely to be sampled near zero, and so forth. We formalize this argument in Section 3.3.2; what is relevant here is that this pattern of poor convergence can be a particular problem when the true group-level variance is near zero, which is common in multilevel models after group-level predictors have been included.

The key parameter expansion step is to include a redundant multiplicative parameter:

$$y_j | \mu, \xi_j \stackrel{\text{indep.}}{\sim} N(\mu + \alpha \xi_j, \sigma_j^2), \quad \text{with } \xi_j \stackrel{\text{iid.}}{\sim} N(0, \sigma_\xi^2), \quad (1.10)$$

where the regression parameters of interest are now $\beta_j = \alpha \xi_j$, with group-level standard deviation $\sigma_\beta = |\alpha| \sigma_\xi$. As has been shown by van Dyk and Meng (2001) and Liu (2003), and also by Liu, Rubin, and Wu (1998) in the context of the EM algorithm, computation of model (1.10) is straightforward.

The likelihood function, $p(y_j | \mu, \sigma_\beta)$, obtained by integrating out the ξ_j 's, is the same as that obtained by integrating out the β_j 's under (1.9). Since we do not alter the prior distribution on (μ, σ_β) , we can fit either model to obtain a Monte Carlo sample from the same posterior distribution $p(\mu, \sigma_\beta | y)$. The added parameter α is not necessarily of any statistical interest, but expanding the model in this way can allow faster convergence of the Gibbs sampler.

2. ALGORITHMS USING MARGINAL AND CONDITIONAL AUGMENTATION

2.1 MOTIVATION AND NOTATION FOR THE EXPANDED MODEL

As we discuss in detail in Section 3.3.2, any of the Gibbs samplers discussed in Section 1 can be slow to converge when the estimated hierarchical variance parameters are near zero. The problem is that, if the current draw of $\sigma_{\beta k}$ is near 0, then in the updating step of β , the deviation of the parameters β in batch k from their prior means in β_0 will themselves be estimated to be very close to 0 (because their prior distribution has scale $\sigma_{\beta k}$). Then, in turn, the variance parameter will be estimated to be close to 0 because it is updated based on the relevant β_j 's (see (1.8)). Ultimately, the stochastic nature of the Gibbs sampler allows it to escape this trap but this may require many iterations.

Van Dyk and Meng (2001) showed how the method of marginal augmentation can substantially improve convergence in this setting; see also Meng and van Dyk (1999), Liu, Rubin, and Wu (1998), van Dyk (2000), and Liu (2003). In order to introduce the working

parameter into the standard model given in (1.2), we follow the parameterization of Liu, Rubin, and Wu (1998), which in our notation is

$$A = [\text{Diag}(W_\beta \alpha)]^{-1}, \quad (2.1)$$

where W_β is the matrix of 0's and 1's from (1.3) that indicates the batch corresponding to each β_j , and α is a $(K_\beta \times 1)$ vector of working parameters. Setting $\zeta = A\beta = [\text{Diag}(W_\beta \alpha)]^{-1} \beta$ yields the reformulated model,

$$y|\zeta, \alpha \sim N(XA^{-1}\zeta, \Sigma_y) \quad (2.2)$$

$$\zeta|\alpha \sim N(\zeta_0, \Sigma_\zeta) \quad (2.3)$$

$$\alpha \sim N(\alpha_0, \Sigma_\alpha), \quad (2.4)$$

where $\zeta_0 = A\beta_0$ and $\Sigma_\zeta = A\Sigma_\beta A$. This prior variance matrix is diagonal since both A and Σ_β are diagonal: $\Sigma_\zeta = \text{Diag}(W_\beta(\sigma_\beta/\alpha)^2)$, where the division and squaring are componentwise. In the limit as $\Sigma_\alpha \rightarrow \infty$, the parameter α becomes nonidentified, but the marginal distribution (β, σ_β) is still defined. As we discuss in Section 1.5, care must be taken when constructing samplers under this limiting working prior distribution.

The transformation $\zeta = A\beta$ introduces one parameter α_k for each of the hierarchical variance parameters $\sigma_{\beta k}^2$. We expect samplers constructed using the appropriate conditional distributions of (2.2)–(2.4) to perform better than the corresponding sampler constructed with (1.1)–(1.2). Indeed, we shall see that, with this working parameter, the Gibbs sampler is less prone to move slowly in regions of the parameter space with $\sigma_{\beta k}$ near zero.

Adding the new parameter α potentially generalizes the hierarchical model in an interesting direction—allowing interactions among the parameters (which is different than interactions among the regression predictions); see Gelman (2004). Our primary purpose in this article, however, is to use the expanded model to improve computational efficiency.

2.2 EQUIVALENCE BETWEEN THE PARAMETER-EXPANDED MODEL AND THE USUAL MODEL

The expanded model can be viewed as a generalization of the usual hierarchical regression model on β , expanding the class of prior distributions on the variance parameter σ_β (as described by Gelman (2006) for the one-variance-parameter case). When the prior distribution on the additional parameter α is proper, one can construct a Gibbs sampler on the expanded space and marginalize by simply focusing on the inferences for the β, σ_β . When α is assigned an improper prior distribution, it will also have an improper posterior distribution, and we have to be more careful in defining a Gibbs sampler with the appropriate marginal distribution on the identified parameter.

The sampling distribution $p(y|\sigma)$ implied by (2.2)–(2.4) is the same as that implied by (1.1)–(1.2) for any *fixed* value of α or if we average over any *proper* working prior distribution on α . In this sense the models are equivalent and α meets the definition of a working parameter. When drawing inferences under model (2.2)–(2.4), however, we must account for the parameters' transformation:

Model (1.1)–(1.2)	Model (2.2)–(2.4)
the vector β	the vector $(W_\beta \alpha) * \zeta$
a single β_j	$\alpha_{k(j)} \zeta_j$
the vector σ_β	the vector $ \alpha * \sigma_\zeta$
a single $\sigma_{\beta k}$	$ \alpha_k \sigma_{\zeta k}$,

where $*$ represents componentwise multiplication.

Thus, if we would like to work with model (1.1)–(1.2), but we use model (2.2)–(2.4) for computational efficiency, then the quantities on the right side of the above table should be reported. For each batch k of regression coefficients, the coefficients ζ_j and the variance parameters $\sigma_{\zeta k}$ sampled under model (2.2)–(2.4) must be multiplied by the corresponding sampled scale parameter α_k so that they can be given their proper interpretations as β_j 's and $\sigma_{\beta k}$ in the original model (1.1)–(1.2).

In fact, with these prior distributions the parameters α_k and $\sigma_{\zeta k}$ cannot be separately identified and only affect the final inferences under the model through their product. If proper prior distributions are assigned, then α and σ_ζ can be interpreted separately as part of a multiplicative-parameter model (see Gelman 2004, sec. 5.2).

2.3 GIBBS SAMPLER COMPUTATION FOR THE EXTENDED MODEL

Under a proper working prior distribution, there are many possible implementations of the Gibbs sampler under the expanded model. For example, we can iteratively sample $(\alpha, \sigma_y, \sigma_\zeta)$ and ζ from their respective complete conditional distributions. Given ζ , we can express (2.2) as

$$y \sim N(X \text{Diag}(\zeta) W_\beta \alpha, \Sigma_y). \quad (2.5)$$

This is a normal linear regression of y on the columns of the known design matrix, $X \text{Diag}(\zeta) W_\beta$ with regression coefficient α . In our model as presented with a diagonal data variance matrix (and more generally, with sufficient structure on Σ_y) we can sample σ_y from $p(\sigma_y | \zeta, y)$. (Likewise, we can independently sample σ_ζ using (2.4).) Given Σ_y , a conditional posterior simulation of α can be obtained by running a standard regression program (augmenting the data vector, design matrix, and the variance matrix to include the Gaussian prior distribution) to obtain an estimate $\hat{\alpha}$ and covariance matrix, then drawing from the multivariate normal distribution centered at that estimate and with that covariance matrix (see, e.g., Gelman et al. 1995). The multiplicative parameter α is typically a relatively short vector, and so we are not bothered by simply updating it in vector fashion using linear regression computations.

Similarly, to update ζ , we express (2.2) as

$$y \sim N(X \text{Diag}(W_\beta \alpha) \zeta, \Sigma_y) \quad (2.6)$$

and work with a regression in which $X \text{Diag}(W_\beta \alpha)$ is the design matrix.

2.4 CONDITIONAL AUGMENTATION IN ONE-AT-A-TIME ALGORITHMS

Rather than sampling α (or equivalently, A) in the iteration, the method of conditional augmentation aims to fix the working parameter at some optimal or nearly optimal value in

terms of the convergence properties of the resulting sampler. For conditional augmentation, rather than sampling $A = [\text{Diag}(W_\beta \alpha)]^{-1}$, we simply allow it to be any fixed matrix. Thus, α amounts to an index on a class of transformations, and we can simply consider ζ to be a transformation of β .

Suppose we do not sample A in the iteration, but draw the variance components jointly given ζ and y and draw the components of ζ one at a time given the variance components and y . If we take A to be an identity matrix, we recover the one-at-a-time Gibbs sampler, so this formulation constitutes a generalization of that in Section 1.3. Since we choose A to improve computational performance, one possibility is to set $A = [R(\sigma^{(t)})]^{-1}$, where $\sigma^{(t)}$ represents the current draw of the variance parameters and R comes from the QR decomposition of $\Sigma_*^{-1/2} X_*$ (see (1.6)) at each iteration, which orthogonalizes the elements of ζ and results in an algorithm that is equivalent to the all-at-once updating. Unfortunately, this strategy does not save time because it requires the computation of $\hat{\beta}$ and R —that is, the full regression computation—at each iteration.

A less computationally burdensome method fixes A at some nearly optimal value that is constant across iterations and thus does not need to be recomputed. A natural approach is to set $A = [R(\hat{\sigma}_0)]^{-1}$, where $\hat{\sigma}_0$ is some estimate of the variance parameters, perhaps estimated by running a few steps of an EM-type algorithm (e.g., van Dyk 2000) or by using an initial run of the Gibbs sampler. Given $\hat{\sigma}_0$, we can run the regression just once, to compute A , and then run the Gibbs sampler conditional on A , that is, using ζ . As long as $R(\sigma)$ is reasonably stable, using a reasonable estimate of σ should keep the components of ζ close to independent in their conditional posterior distribution. (We expect this to typically be a useful strategy since the design matrix X and the fixed structure of the covariance matrices Σ_* do not change as the simulation progresses, and this induces stability in R .)

As with structured MCMC, we aim to develop a sampler with the low autocorrelation of the all-at-once Gibbs sampler but with the quick per-iteration computation time of the one-at-a-time Gibbs sampler. Also as with structured MCMC, our strategy involves an approximation to the conditional posterior variance matrix of the regression coefficients. An advantage of our proposal, however, is that it does not require a Metropolis–Hastings correction to adjust for the approximation. The relative computational performance of the two strategies, nonetheless, depends on the quality of the approximations.

A natural modification is to set A to the posterior mean of R , which can be computed, for example, by updating occasionally (e.g., every 200 steps) online; for example, set $A^{(0)} = R(\hat{\sigma}_0)^{-1}$ and then, for $t = 200, 400, \dots$,

$$A^{(t)} = \frac{1}{t + 200} \left[{}_t A^{(t-200)} + 200 [R(\sigma^{(t)})]^{-1} \right].$$

This updating would take place only during the burn-in period (typically half the total length of the chain; see Gelman and Rubin 1992) so as not to interfere with the Markov property of the ultimate chain.

2.5 SETTING UP THE COMPUTATION

We perform all computations in terms of the regression model (2.2)–(2.4), without assuming any special structure in the design matrix X . The algorithms must begin with estimates of the variance parameters σ and also, for the one-at-a-time algorithms, an estimate of ζ . In order to reliably monitor the convergence of the Gibbs sampler, it is desirable to start the Gibbs sampler runs at a set of initial values that are “overdispersed,” that is, from a distribution that includes and is more variable than the target distribution. More precisely, it is desired that the simulation draws from the mixture of all the simulated sequences should decrease in variability as the simulation proceeds, with the variability within each sequence increasing, so that approximate convergence can be identified with the empirical mixing of the simulated sequences (see Gelman and Rubin 1992). When using over-dispersed starting values, it becomes particularly important to consider the behavior of the algorithms when they are started far from the posterior mode, as we discuss in Section 3.3.

3. THEORETICAL ARGUMENTS

The previous section introduced augmentation algorithms and suggested why they should be able to speed convergence by reducing dependence among regression coefficients (by a matrix multiplication that is equivalent to rotating the parameter space) and reducing dependence between batches of coefficients and their variance parameters (using multiplicative redundant parameters). After a brief look at computation time, we provide theoretical arguments to show the potential gains from these ideas in Gibbs sampling. We work out why parameter expansion works well in a one-way hierarchical model that is a simple case relative to the generality to which the methods are applicable; we believe these theoretical arguments provide intuition about why they should work in more complex examples. Our most detailed treatment is of the convergence of the expanded algorithm in Section 3.3, because this is where we are providing a novel discussion of the algorithm’s ability to avoid being stuck near the boundary of the parameter space.

3.1 COMPUTATION TIME PER ITERATION

The QR decomposition of the W matrix and backsolving the two upper-triangular systems that are required for the vector updating algorithms take $O(2nm^2)$ floating-point operations (flops), where n and m are the length of y and ζ , respectively; see Golub and van Loan (1983). For the scalar updating algorithm, updating all m components of ζ and transforming back to ζ takes only $O(10nm)$ flops. The computation for updating α require another $O(nm)$ flops. (We assume that the length of α and the number of variance parameters is negligible compared to m and n .) In many of the problems we work with, m is quite large—typically some fraction of n —and thus the scalar updating algorithms require $O(n^2)$ flops per iteration, compared to $O(n^3)$ for the vector algorithms.

3.2 CONVERGENCE RATE OF VECTOR VS. SCALAR UPDATING OF REGRESSION PARAMETERS

An interesting tradeoff occurs as the number of predictors becomes large. As the batch sizes $n_{\beta k}$ increase, the dimensions of the X_* matrix in (1.5) also increase, making the least-squares regression computation slower. However, having more replication increases the precision of the estimates of the variance parameters, which means that once they have been reasonably estimated, a one-time transformation as described in Section 2.4 will do a good job of making the components of ζ close to independent. Thus, *increasing the number of regression coefficients and the number of parameters per batch (the $n_{\beta k}$'s) can make the Gibbs sampler computations more stable on the transformed space.*

We shall try to understand the efficiency of the vector and scalar algorithms by considering different amounts of knowledge about the variance parameters σ in the model (see (1.3)). In this discussion we focus on the rotation-to-approximate-independence described in Section 2.4. Thus ζ refers to the transformations of β described in Section 2.4.

We can understand the efficiencies of vector and scalar updating by considering three scenarios involving inference about the variance parameters.

First suppose all the variance parameters are known. Then the vector updating of ζ converges in a single step, meaning that the Gibbs sampler draws are equivalent to independent draws from the posterior distribution of ζ . In contrast, the speed of the untransformed scalar updating algorithm is roughly determined by the second eigenvalue of the posterior variance matrix V_ζ . In many hierarchical models, it is possible to keep the correlations in V_ζ low by centering and scaling the parameter ζ (see Gelfand, Sahu, and Carlin 1995; Gilks and Roberts 1996), but in general finding a good transformation can require matrix operations that can be slow in high dimensions. Without such a rotation, the computational savings from the scalar updating algorithm may be lost if they require many more iterations to converge. If all variance parameters are known, the transformed scalar updating algorithm will converge in one step, and be just as efficient as the vector algorithm.

Second, consider a setting where the variance parameters are unknown but are accurately determined by the posterior distribution. This should be the case in large datasets with many observations per group and many groups. In this case, the transformed scalar updating should be nearly as efficient as the vector algorithm, since the only reason the orthogonalizing transformation varies is uncertainty in Σ_y and Σ_ζ .

Finally, suppose that the variance parameters are poorly estimated, as could occur if some of the variance components $\sigma_{\zeta k}$ had few associated ζ_j 's, along with a weak or non-informative prior distribution. In this case, the transformed scalar updating, with any particular transformation, might not work very well, and it would make sense to occasionally update the transformation during the burn-in period based on current values of Σ_y and Σ_ζ in the simulation, as discussed in Section 2.4.

3.3 EFFECT OF PARAMETER EXPANSION ON CONVERGENCE RATE

Because the parameter expansion operates independently for each hierarchical variance component k , we consider its effect on the convergence of these components separately.

Ideally, the marginal augmentations methods can outperform the standard Gibbs sampler in three settings:

- $\sigma_{\beta k}$ near the posterior mode (relevant in problems for which $\sigma_{\beta k}$ is well estimated in the posterior distribution);
- $\sigma_{\beta k}$ near 0 (relevant because of possible extreme starting points and also for problems in which $\sigma_{\beta k}$ has a nontrivial posterior probability of being near 0); and
- $\sigma_{\beta k}$ near ∞ (relevant because of possible extreme starting points and also for problems in which $\sigma_{\beta k}$ has a non-trivial posterior probability of being large).

For each of these cases, we consider first the speed of the corresponding EM algorithms and then consider the Gibbs sampler. Because EM algorithms are deterministic, computing their speed of convergence is easier than for Gibbs samplers. Due to the similarity in the structure of the two classes of algorithms, however, the convergence properties of EM algorithms can emulate those of Gibbs samplers (e.g., van Dyk and Meng 2001). The EM algorithms we shall consider are those that treat the components of ζ as missing data and thus converge to the marginal posterior mode of $(\alpha, \sigma_y, \sigma_\zeta)$, averaging over ζ .

We work through the details in the context of a simple model with one variance component, $\sigma_\beta = \alpha\sigma_\zeta$, in order to illustrate the principles that we conjecture to hold for hierarchical models in general. Consider the following model, with variance expressed in parameter-expanded form:

$$\begin{aligned} \text{For } j = 1, \dots, m: \quad y_j | \zeta_j &\sim \text{N}(\alpha\zeta_j, 1) \\ \zeta_j &\sim \text{N}(0, \sigma_\zeta^2) \\ p(\alpha, \sigma_\zeta) &\propto 1. \end{aligned}$$

The Gibbs sampler for this model is

$$\begin{aligned} \zeta_j &\leftarrow \text{N}\left(\frac{1}{\alpha} \frac{y_j}{1 + \frac{1}{\sigma_\beta^2}}, \frac{1}{\alpha^2} \frac{1}{1 + \frac{1}{\sigma_\beta^2}}\right), \quad \text{for } j = 1, \dots, m \\ \alpha &\leftarrow \text{N}\left(\frac{\sum_{j=1}^m \zeta_j y_j}{\sum_{j=1}^m \zeta_j^2}, \frac{1}{\sum_{j=1}^m \zeta_j^2}\right) \\ \sigma_\zeta^2 &\leftarrow \sum_{j=1}^m \zeta_j^2 / \chi_{m-1}^2. \end{aligned} \tag{3.1}$$

(In this case, the vector and the componentwise algorithms are identical since the ζ 's are independent conditional on the hyperparameters.) The EM updating is simply,

$$\begin{aligned} \alpha &\leftarrow \alpha \frac{s_y^2}{1 + \frac{1}{\sigma_\beta^2} s_y^2 + 1} \\ \sigma_\zeta^2 &\leftarrow \frac{1}{\alpha^2} \left(\frac{1}{1 + \frac{1}{\sigma_\beta^2}} \left(\frac{1}{1 + \frac{1}{\sigma_\beta^2}} s_y^2 + 1 \right) \right), \end{aligned} \tag{3.2}$$

where

$$s_y^2 = \frac{1}{m} \sum_{j=1}^m y_j^2.$$

At each step, σ_β is set to $|\alpha|\sigma_\xi$. The EM algorithm converges approximately to $\hat{\sigma}_\beta = \sqrt{s_y^2 - 1}$, or 0 if $s_y \leq 1$.

3.3.1 Speed of the Algorithms Near the Posterior Mode

For $\sigma_{\beta k}$ near the posterior mode (assuming that mode is not 0), the speed of the two-step Gibbs sampler is roughly the same as the corresponding EM algorithm (Liu 1994), with convergence rate depending on the covariance matrix of the approximate Gaussian distribution. Liu, Rubin, and Wu (1997) considered the EM algorithm for the hierarchical normal model and find that parameter expansion is uniformly better than the usual algorithm (which, in the parameter expansion context, corresponds to holding the parameters α_k fixed at 1).

3.3.2 Speed of the Algorithms for σ_β Near 0

For $\sigma_{\beta k}$ near 0, the usual algorithm, in both the EM and Gibbs contexts, is notoriously slow, but we shall see that the parameter expanded versions move much faster.

We first consider the EM algorithm. Under the usual parameterization, α is fixed at 1, and only $\sigma_\beta = \sigma_\xi$ is updated. For σ_β near 0, we can express (3.2) as

$$\begin{aligned} \sigma_\beta &\leftarrow \frac{\sigma_\beta}{1 + \sigma_\beta^2} \sqrt{1 + \sigma_\beta^2 + \sigma_\beta^2 s_y^2} \\ &= \sigma_\beta \left(1 + \frac{1}{2}(s_y^2 - 1)\sigma_\beta^2 + \dots \right). \end{aligned}$$

Thus, even the *relative* change in σ_β at each step approaches 0 as $\sigma_\beta \rightarrow 0$. If $s_y^2 \leq 1$, this means that σ_β will approach 0 at a slower-than-linear rate; if $s_y^2 > 1$ and σ_β happens to be started near 0, then it will move away from 0 hopelessly slowly.

In contrast, the parameter-expanded EM algorithm updates both α and σ_ξ , yielding, for σ_β near 0,

$$\begin{aligned} \sigma_\beta &\leftarrow \sigma_\beta \frac{s_y^2}{\sqrt{1 + (s_y^2 + 1)\sigma_\beta^2}} \\ &= \sigma_\beta (s_y^2 + \dots), \end{aligned}$$

which is linearly convergent, which should be acceptable in practice as long as σ_β is not started at an extremely small value.

We now consider the Gibbs sampler which, under the conventional parameterization, adds two sorts of variation to the EM algorithm: (1) $E(\sum_{j=1}^m \xi_j^2)$ is replaced by the random variable $\sum_{j=1}^m \xi_j^2$, and (2) division by a χ_m^2 random variable in the update of σ_β^2 . (Under

the conventional parameterization, the updating of α in (3.2) does not occur since α is fixed at 1.) When the current value of σ_β in the Gibbs sampler is near 0, each of these steps adds a coefficient of variation $1/(\sqrt{2}m)$ to the updating step for σ_β^2 ; so combined they give the random updating of σ_β a coefficient of variation of $1/m$. Within the context of the usual parameterization, with σ_β near 0, this variation acts like a random walk, and it implies that the Gibbs sampler will require on the order of $m(\log(\sigma_{\beta 0}))^2$ iterations to “escape” from a starting $\sigma_{\beta 0}$ near 0. (See Rosenthal (1995) for more formal versions of this argument.)

For the parameter-expanded algorithm, the Gibbs sampler from (3.1) adds another random component through the updating of α . The updated $\sigma_\beta = |\alpha|\sigma_\xi$ is essentially a least-squares regression coefficient plus noise (the Gibbs update for α), divided by the square root of a χ_{n-1}^2 random variable (the Gibbs update for σ_ξ). This added random component causes the draw to be less dependent on the previous iteration. The updated σ_β can then be written as

$$\sigma_\beta \leftarrow \left| \frac{\sum_{j=1}^m \gamma_j y_j}{\sqrt{\sum_{j=1}^m \gamma_j^2}} + z \right| / x, \quad (3.3)$$

where the components of $\gamma = \xi/\alpha$ can be written as

$$\gamma_j = \frac{1}{1 + \frac{1}{\sigma_\beta^2}} + \frac{1}{\sqrt{1 + \frac{1}{\sigma_\beta^2}}} z_j,$$

and the following random variables are independent:

$$z \sim N(0, 1), \quad x^2 \sim \chi_{m-1}^2, \quad z_j \sim N(0, 1), \quad \text{for } j = 1, \dots, m. \quad (3.4)$$

The random variables in (3.3) induce a distribution for σ_β on the left of (3.3) that depends only on m , $s_y^2 = \frac{1}{m} \sum_{j=1}^m y_j^2$, and the previous value of σ_β . For σ_β near 0, this distribution has a standard deviation on the order of $1/\sqrt{m}$ on the absolute scale. Therefore, it is impossible for σ_β to get stuck at values less than $1/\sqrt{m}$, no matter how small its value was in the previous iteration. This is a distinct improvement over the Gibbs sampler for the conventional parameterization.

This jumpy behavior of the parameter-expanded algorithm near $\sigma_\beta = 0$ is reminiscent of the nonzero lowest energy state in quantum mechanics, which is related to the uncertainty principle—if σ_β is precisely localized near 0, then the “momentum” of the system will be high (in that σ_β is likely to jump far away). When m is large, the minimal “energy” of $1/\sqrt{m}$ becomes closer to 0, which corresponds in a physical system to the classical limit with many particles. In a statistical sense, this behavior is reasonable because the precision of the marginal posterior distribution for σ_β is determined by the number of coefficients m in the batch.

3.3.3 Speed of the Algorithms for σ_β Near ∞

If σ_β is started at a very high value, both the usual parameterization and the parameter-expanded algorithm perform reasonably well. From $\sigma_\beta = \infty$, the usual EM algorithm

moves in one step to $\sigma_\beta = \sqrt{s_y^2 + 1}$ and the parameter-expanded EM algorithm moves in one step to $\sigma_\beta = s_y^2 / \sqrt{s_y^2 + 1}$. Of these two, the parameter-expanded version is preferable (since it is closer to the convergence point, $\sqrt{s_y^2 - 1}$), but they are both reasonable. Similarly, σ_β will not get stuck near ∞ under either Gibbs sampler implementation.

4. EXAMPLES

We consider two examples that have arisen in applied research. For each example and each algorithm, we compute computation time per iteration, number of iterations required for convergence (as determined by the condition that the variance ratio \hat{R} (Gelman and Rubin 1992) is less than 1.2 for all model parameters), and simulation efficiency (measured by the autocorrelations).

4.1 EXAMPLES AND ALGORITHMS

Our first example is a hierarchical model for effects of an educational testing experiment in eight schools, described by Rubin (1981) and Gelman et al. (1995). For each school $j = 1, \dots, 8$, an experiment was performed to estimate the treatment effect β_j in that school; a regression analysis applied to the data from that school alone yielded an estimate of the treatment effect and its standard error, which we label y_j and σ_j , respectively. We model the data using the two-level Gaussian model (1.9). The sample size within each school is large enough to justify the Gaussian assumption in level one of (1.9). We use a noninformative uniform hyperprior density, $p(\mu, \sigma_\beta) \propto 1$ (see Gelman (2006) for discussion of alternative prior distributions). Interest lies in the individual school parameters β_j and also in the hyperparameters (μ, σ_β) ; we label the set of unknown parameters $\gamma = (\beta, \mu, \sigma_\beta)$. This particular example is quite small and should be easy to compute; in fact, since there is only one hierarchical variance component, it is in fact possible to draw posterior simulations noniteratively by first computing the marginal posterior density for σ_β at a grid of points and then simulating the parameters in the order σ_β, μ, β (Rubin 1981).

We consider two standard Gibbs samplers that can be constructed to sample the posterior distribution under this model. The first updates the mean parameters jointly, or as a vector; we refer to this sampler as the V-SAMPLER.

V-SAMPLER:

Step 1: Sample $(\mu^{(t+1)}, \beta^{(t+1)})$ jointly from $p(\mu, \beta \mid \sigma_\beta = \sigma_\beta^{(t)}, y)$.

Step 2: Sample $\sigma_\beta^{(t+1)}$ from $p(\sigma_\beta \mid \mu = \mu^{(t+1)}, \beta = \beta^{(t+1)}, y)$.

A second sampler, updates each of the mean components separately from their respective complete conditional distributions; we call this sampler the S-SAMPLER.

S-SAMPLER:

Step 1: Sample $\mu^{(t+1)}$ from $p\left(\mu \mid \beta = \beta^{(t)}, \sigma_\beta = \sigma_\beta^{(t)}, y\right)$.

Step 2: Sample $\beta_j^{(t+1)}$ from $p\left(\beta_j \mid \mu = \mu^{(t+1)}, \beta_{-j} = \beta_{-j}^{(t+1)}, \sigma_\beta = \sigma_\beta^{(t)}, y\right)$, for $j = 1, \dots, J$, where $\beta_{-j}^{(t+1)} = \left(\beta_1^{(t+1)}, \dots, \beta_{j-1}^{(t+1)}, \beta_{j+1}^{(t)}, \dots, \beta_J^{(t)}\right)$

Step 3: Sample $\sigma_\beta^{(t+1)}$ from $p\left(\sigma_\beta \mid \mu = \mu^{(t+1)}, \beta = \beta^{(t+1)}, y\right)$.

A possible difficulty for the convergence of both of these Gibbs samplers occurs when the maximum likelihood estimate of the hierarchical scalar parameter, σ_β , is zero. There is also a substantial portion of the posterior density near $\sigma_\beta = 0$, and there is the possibility that the algorithm may move slowly in this region.

To improve computational efficiency, we consider two marginal Gibbs samplers that are based on a working parameter α that is introduced via $\zeta = \beta/\alpha$. Introducing this transformation into (1.9), the model becomes (1.10), where $\sigma_\zeta = \sigma_\beta/|\alpha|$. We can fit either model (1.9) or (1.10) to obtain a Monte Carlo sample from the same posterior distribution $p(\mu, \sigma_\beta|y)$. In the Appendix we further show that with the improper prior distribution specified below on α , all the samplers return a sample from the same target joint posterior distribution $p(\beta, \mu, \sigma_\beta|y)$.

To derive the samplers under model (1.10), we begin with the proper working prior distribution, $\alpha \sim \text{Inv-gamma}(\xi_0, \xi_1)$, which implies $p(\mu, \sigma_\zeta^2, \alpha) \propto \alpha^{-\xi_0} \exp(-\xi_1/\alpha)/\sigma_\zeta^2$. We then construct two Gibbs samplers on the joint proper posterior distribution $p(\mu, \zeta, \sigma_\zeta, \alpha|y)$; the two samplers use updating schemes that correspond to the V-SAMPLER and the S-SAMPLER, respectively. The chains are constructed so that the marginal chains $\{\gamma^{(t)}, t = 1, 2, \dots\}$ with $\gamma = (\mu, \beta, \sigma_\beta)$ are themselves Markovian. We then take the limit of the transition kernels for the two chains as the working prior distribution becomes improper. We call these limiting kernels the V+PX-SAMPLER and the S+PX-SAMPLER, respectively. Here we state these limiting samplers; we prove that their stationary distributions are both the target posterior distribution, $p(\gamma|y)$ in the Appendix.

V+PX-SAMPLER:

Step 1: Sample $(\mu^{(t+1)}, \beta^*)$ jointly from $p\left(\mu, \beta \mid \sigma_\beta = \sigma_\beta^{(t)}, y\right)$; this is the same distribution as is sampled in Step 1 of the V-SAMPLER.

Step 2: Sample σ_β^* from $p\left(\sigma_\beta \mid \mu = \mu^{(t+1)}, \beta = \beta^*, y\right)$; this is the same distribution as is sampled in Step 2 of the V-SAMPLER.

Step 3: Sample α from $p\left(\alpha \mid \mu = \mu^{(t+1)}, \zeta = \beta^*, \sigma_\zeta = \sigma_\beta^*, y\right)$; that is, sample

$$\alpha \sim \text{N} \left[\frac{\sum_{j=1}^J \beta_j^* (y_j - \mu^{(t+1)}) / \sigma_j^2}{\sum_{j=1}^J (\beta_j^*)^2 / \sigma_j^2}, \left(\sum_{j=1}^J (\beta_j^*)^2 / \sigma_j^2 \right) \right]. \quad (4.1)$$

Step 4: Compute $\beta^{(t+1)} = \alpha\beta^*$ and $\sigma_\beta^{(t+1)} = |\alpha|\sigma_\beta^*$.

We use a star in the superscript to represent intermediate quantities that are not part of the transition kernel.

The S+PX-SAMPLER begins with Step 1 and Step 2 of the S-SAMPLER, but records what is recorded as $\beta^{(t+1)}$ and $\sigma_\beta^{(t+1)}$ in the S-SAMPLER as β^* and σ_β^* , respectively. The iteration is completed with Step 3 and Step 4 of the V+PX-SAMPLER.

In the second example, we consider the forecast skill of three global circulation models (GCM) based on Africa precipitation data in the fall (October, November, December) season. The models divide the globe into a grid, on which Africa covers 527 boxes, and we have 41 observed precipitation values (between 1950 and 1990) for each box in a given season. Here we consider three GCM models, each of which gives us 10 predicted precipitation values for each box in a given season (Mason et al. 1999; Rajagopalan et al. 2000). In this example, we let y_{jt} and $x_{jt}^{m,k}$ represent observed and the k th predicted precipitation anomaly for the m th GCM ensemble (out of $M = 3$ models) in box j at time t . We use as predictors the average values,

$$\bar{x}_{jt}^m = \frac{1}{10} \sum_{k=1}^{10} x_{jt}^{m,k}.$$

We set up the following multilevel model using y_{jt} as the response and \bar{x}_{jt}^m as predictors in a Bayesian framework:

$$y_{jt} = \beta_j + \delta_t + \sum_{m=1}^M \beta_j^m \bar{x}_{jt}^m + \epsilon_{jt}, \quad \text{for } j = 1, \dots, 527 \text{ and } t = 1, \dots, 41. \quad (4.2)$$

The parameters in the model are defined as follows:

- δ_t is an offset for time t ; we assign it a normal population distribution with mean 0 and standard deviation σ_δ .
- β_j is an offset for location j ; we assign it a normal population distribution with mean μ_β and standard deviation σ_β .
- β_j^m is the coefficient of ensemble forecast m in location j ; we assign it a normal population distribution with mean μ_{β_m} and standard deviation σ_{β_m} .
- ϵ_{jt} 's are independent error terms assumed normally distributed with mean 0 and standard deviation σ_y .

In this example, our parameter expansion model is

$$y_{jt} = \alpha_\beta \tilde{\beta}_j + \alpha_\delta \tilde{\delta}_t + \sum_{m=1}^M \alpha_{\beta m} \tilde{\beta}_j^m \bar{x}_{jt}^m + \epsilon_{jt}, \quad \text{for } j = 1, \dots, 527 \text{ and } t = 1, \dots, 41, \quad (4.3)$$

using the notation $\tilde{\beta}$, and so on, for unscaled parameters (so that $\beta_j = \alpha_\beta \tilde{\beta}_j$ for each j ; $\delta_t = \alpha_\delta \tilde{\delta}_t$ for each t ; and so forth). We assign uniform prior distributions on the (unidentifiable) multiplicative parameters α and the mean and scale hyperparameters for the prior




Figure 1. Computation time of the four algorithms for the eight schools example (plotted on a logarithmic scale). V: vector updating, S: scalar updating, V+PX: vector updating with parameter expansion, and S+PX: scalar updating with parameter expansion. The dots display the combination of the computation time per iteration and the average number of iterations required for approximate convergence of each algorithm. The lines are indifference curves in total computation time for convergence to the stationary distribution.

distributions on the $\tilde{\beta}$'s, $\tilde{\delta}$'s, and so forth. We compute the samplers for the expanded models and then save the simulations of the parameters as defined in (4.2). For this problem it was most direct to simply set up the model in the expanded parameter space.

4.2 RESULTS

We present the results of the standard and PX-Gibbs algorithms. For each algorithm, starting points were obtained by running the EM algorithm (with initial guesses of 1 for all the variance components) and then drawing from a t_4 distribution. When the EM estimate of a variance component was zero (as in the study of the eight schools), we used 1 as a starting point.

Figure 1 compares the computation time of the standard and PX-Gibbs algorithms for the example of the eight schools. Each time we run 10 chains until approximate convergence ($\hat{R} < 1.2$ for all parameters). In Figure 1, V represents vector updating without parameter expansion, S represents scalar updating without parameter expansion, V+PX represents vector updating with parameter expansion, and S+PX represents scalar updating with parameter expansion. The dots display the combination of the average computation time per iteration and the average number of iterations required for convergence for each algorithm. The lines represent indifference curves for total computation time until convergence to the target distribution. These are straight lines because the graph is on the logarithm scale. A point on a lower line indicates a more efficient algorithm in total computation time. Therefore the most efficient algorithm is scalar updating with parameter expansion, which only takes an average of 0.39 seconds in total computation time per




Figure 2. Simulation efficiency (measured by the autocorrelations) of variance component σ_β for the eight schools example. The sum of the absolute autocorrelation functions (ACF) are given and are roughly equal to the expected sample sizes (ESS).

Gibbs sampler chain; the second is the vector updating with parameter expansion, which takes 0.69 seconds; the third is scalar updating, which takes 4.2 seconds; and the slowest is vector updating, which takes 8.7 seconds. In this example, parameter expansion is 22 times faster than the traditional algorithms.

The Gibbs samplers on the original scale (points V and S in Figure 1) are slow to converge because the group-level variance parameter, σ_β , is estimated to be close to zero in this example. In fact, the marginal posterior mode of σ_β is zero; thus the Gibbs sampler spends a lot of time down there and can easily get stuck in the models without parameter expansion.

Figure 2 shows the simulation efficiency (measured by the autocorrelations and the sum of their absolute values) for the variance component σ_β . The vector and scalar updating with parameter expansion are much more efficient than the other two algorithms.

Because the design matrix for the second example (climate modeling) is large, it is difficult to implement vector updating in R or S-Plus. Although sparse matrix methods might help with vector updating, for the purposes of illustration we consider only the scalar updating algorithm. The computation speed per iteration for scalar updating in our simulation was similar with or without parameter expansion. However, the scalar updating with parameter expansion needed only 400 iterations to reach convergence, compared to 10,000 for the scalar updating without parameter expansion. Figure 3 compares the computation time of the two algorithms for the second example (climate modeling). The most efficient algorithm is scalar updating with parameter expansion, which takes an average of 500 sec-




Figure 3. Computation time (on a logarithmic scale) of the two algorithms for the climate modeling example. S: scalar updating and S+PX: scalar updating with parameter expansion. The dots display the combination of the computation time per iteration and the iterations required for convergence for each algorithm. The lines are indifference curves in total computation time.

onds per chain in total computation time compared to 13,000 seconds per chain for scalar updating.

5. GENERALIZED LINEAR MODELS

In this section we discuss how the methods developed in Sections 2 and 3 for hierarchical linear models can be extended to hierarchical generalized linear models, so that (1.1) is replaced by

$$y|\beta \sim \text{glm}(X\beta) \quad \text{or} \quad y|\beta \sim \text{glm}(X\beta, \Sigma_y); \quad (5.1)$$

the simpler form is used for models such as the binomial and Poisson with no free variance parameters. We use the general “glm” notation to include the necessary probability models and link functions (McCullagh and Nelder 1989).

5.1 A METROPOLIS ADAPTATION TO THE SIMPLE GIBBS SAMPLERS

For a hierarchical generalized linear model, the simple all-at-once and one-at-a-time Gibbs samplers described in Section 1.3 must be modified since the likelihood is not conjugate with the Gaussian prior distribution. The most straightforward adaptation is to perform an approximate Gibbs sampler, drawing from the conditional posterior distribution corresponding to the approximate normal likelihood and then accepting or rejecting at each step based on the Metropolis–Hastings rule. For the one-at-time algorithm, one approach for the one-dimensional jumps is the adaptive Metropolis algorithm of Gilks, Best, and Tan (1995). For the all-at-once algorithm, a natural approach to updating β is, by analogy to

maximum likelihood computations, to run the regression based on a linearization of the likelihood at the current values of the variance components. This involves computing the conditional (on the variance components) posterior mode and second derivative matrix and using a multivariate Gaussian or t distribution to generate proposals.

5.2 PARAMETER EXPANSION FOR GENERALIZED LINEAR MODELS

We can use marginal augmentation for the generalized linear model if we simply replace (2.2) by

$$y|\zeta, \alpha \sim \text{glm}(X((W_\zeta \alpha) * \zeta)) \quad \text{or} \quad |\zeta, \alpha \sim \text{glm}(X((W_\zeta \alpha) * \zeta), \Sigma_y),$$

with the equivalences to the original model as described in Section 2.2. As in Section 2.3 we use a proper working prior distribution and again the Gibbs sampler for the new model must include a step to update α . Since the model is nonconjugate, this can be performed by Metropolis jumping, or via a linearization of the model $y \sim \text{glm}(X((W_\zeta \alpha) * \zeta), \Sigma_y)$ —considered as a likelihood for α —followed by a draw from the corresponding approximate Gaussian conditional prior distribution for α and a Metropolis–Hastings accept/reject step. Computation for ζ can similarly be performed using a Metropolis jump or a Metropolis–approximate–Gibbs. In either case, we want an approximate transformation to independence (as in Section 2.4), whether for scaling the Metropolis proposal distribution or approximating the Gibbs sampler. Finally, the variance parameters can be updated using the Gibbs sampler as with the normal model, since they are linked to the other model parameters through the prior distribution, not the likelihood.

5.3 ADAPTATION OF THE ONE-AT-A-TIME ALGORITHMS

If the components of β are highly correlated in their posterior distribution, then the Metropolis–Hastings sampler corresponding to the one-at-a-time Gibbs sampler can move slowly. To improve the sampler, we can adapt the rotation method based on an approximation to the posterior covariance matrix of the regression parameters. In particular, we can apply Metropolis–Hastings algorithms to the components of ζ one at a time, either as corrections to approximate Gibbs sampler jumps for generalized linear models (where the nonconjugate conditional posterior densities make exact Gibbs impractical), or simply using spherically symmetric Metropolis jumps on ζ , starting with a unit normal kernel with scale $2.4/\sqrt{d}$ (where d is the dimension of β) and tuning to get an approximate acceptance rate of 1/4 (see Gelman, Roberts, and Gilks 1996).

6. DISCUSSION

We see this article as having three main contributions. First, we combine some ideas from the recent statistical literature to construct a family of improved algorithms for posterior simulations from hierarchical models. We also suspect that the general ideas of approximate rotation for correlated parameters and parameter expansion for variance components will be useful in more elaborate settings such as multivariate and nonlinear models.

Second, the computations are set up in terms of an expanded model, following the work of Liu, Rubin, and Wu (1997) for the EM algorithm, and more recently called “redundant parameterization” in the context of multilevel models (Gelman and Hill 2007). Once this model has been set up, the next natural step is to see if its additional parameters can be given their own statistical meaning, as discussed in Section 2.3. There is a history in statistics of new models that are inspired by computational developments (Gelman 2004). For example, Besag (1974) motivated conditional autoregression by way of the Hammersley–Clifford theorem for joint probability distributions, and Green (1995) introduced a reversible-jump Markov chain algorithm that has enabled and motivated the use of mixture posterior distributions of varying dimension. Multiplicative parameter expansion for hierarchical variance components is another useful model generalization that was originally motivated for computational reasons (see Gelman 2006).

Third, we connect computational efficiency to the speed at which the various iterative algorithms can move away from corners of parameter space, in particular, near-zero estimates of variance components. When the number of linear parameters m in a batch is high, the corresponding variance component can be accurately estimated from data, which means that a one-time rotation can bring the linear parameters to approximate independence, leading to rapid convergence with one-at-a-time Gibbs or spherical Metropolis algorithms. This is a “blessing of dimensionality” to be balanced against the usual “curse.” On the other hand, the “uncertainty principle” of the parameter-expanded Gibbs sampler keeps variance parameters from being trapped within a radius of approximately $1/\sqrt{m}$ from 0 (see the end of Section 3.3.2), so here it is helpful if m is low.

Further work is needed in several areas. First, it would be good to have a better approach to starting the Gibbs sampler. For large problems, the EM algorithm can be computationally expensive—and it also has the problem of zero estimates. It should be possible to develop a fast and reliable algorithm to find a reasonably over-dispersed starting distribution without having to go through the difficulties of the exact EM algorithm. A second, and related, problem is the point estimate of (σ, σ_β) used to compute the estimated covariance matrix R_0 required for the scalar updating algorithms with transformation. Third, the ideas presented here should be generalizable to multivariate models as arise, for example, in decompositions of the covariance matrix in varying-intercept, varying-slope models (O’Malley and Zaslavsky 2005; MacLehose et al. 2007). Finally, as discussed by van Dyk and Meng (2001) and Liu (2003), the parameter expansion idea appears open-ended, which makes us wonder what further improvements are possible for simple as well as for complex hierarchical models.

A. APPENDIX: VERIFYING THE STATIONARY DISTRIBUTION OF SAMPLERS BASED ON MARGINAL AUGMENTATION

We consider Example 1 of Section 4.1. To prove that $p(\gamma | y)$ is the stationary distribution of both the V+PX-SAMPLER and the S+PX-SAMPLER we use Lemma 2 of Liu and Wu (1999). The lemma states that if a sequence of proper Markovian transition kernels each have the same stationary distribution, and if the sequence converges to a proper

Markovian transition kernel, the limiting kernel has the same stationary distribution as the sequence. We construct the sequence of proper Markovian transition kernels using a sequence of proper prior distributions on α , $p_n(\alpha)$, that converge to the improper prior distribution, $p_\infty(\alpha) \propto 1/\alpha$. Namely, we use $\alpha \sim \text{Inv-gamma}(\omega_n, \omega_n)$ where $\omega_n \rightarrow 0$ as $n \rightarrow \infty$.

Consider the following transition kernel constructed under some proper $p_n(\alpha)$.

PROPER V+PX-SAMPLER

Step 1: Sample $(\mu^{(t+1)}, \zeta^*, \alpha^*)$ from $p_n(\mu, \zeta, \alpha \mid \sigma_\beta = \sigma_\beta^{(t)}, y)$.

Step 2: Sample σ_ξ^* from $p(\sigma_\xi \mid \mu = \mu^{(t+1)}, \zeta = \zeta^*, \alpha = \alpha^*, y)$.

Step 3: Sample $\alpha^{(t+1)}$ from $p_n(\alpha \mid \mu = \mu^{(t+1)}, \zeta = \zeta^*, \sigma_\xi = \sigma_\xi^*, y)$.

Step 4: Set $\beta^{(t+1)} = \alpha^{(t+1)}\zeta^*$ and $\sigma_\beta^{(t+1)} = |\alpha^{(t+1)}|\sigma_\xi^*$.

Here we use the subscript n to emphasize the dependency of certain conditional distributions on the choice of $p_n(\alpha)$. For any proper $p_n(\alpha)$, the stationary distribution of this sampler is $p(\gamma, \alpha|y) = p(\gamma|y)p_n(\alpha)$. Moreover, the marginal chain $\{\gamma^{(t)}, t = 1, 2, \dots\}$ is Markovian with stationary distribution $p(\gamma|y)$ for any proper $p_n(\alpha)$. Thus, in order to establish that the stationary distribution of V+PX-SAMPLER is also $p(\gamma|y)$, we need only show that the limit of the sequence of transition kernels constructed using the PROPER V+PX-SAMPLER is the Markovian kernel given in the V+PX-SAMPLER.

Proof: Step 1 and Step 2 of the PROPER V+PX-SAMPLER can be rewritten as

- Sample α^* from $p_n(\alpha \mid \sigma_\beta^{(t)}, y) = p_n(\alpha)$.

- Sample

$$(\mu^{(t+1)}, \beta^*) \text{ jointly from } p(\mu, \beta \mid \sigma_\beta^{(t)}, y). \quad (\text{A.1})$$

- Sample

$$\sigma_\beta^* \text{ from } p(\sigma_\beta \mid \mu = \mu^{(t+1)}, \beta = \beta^*, y). \quad (\text{A.2})$$

- Set $\zeta^* = \beta^*/\alpha^*$ and $\sigma_\xi^* = \sigma_\beta^*/|\alpha^*|$.

Only the draw of α^* depends on $p_n(\alpha)$.

To analyze Step 3 in the limit, we note that because $p_n(\alpha) \rightarrow p_\infty(\alpha)$, $p_n(\mu, \zeta, \sigma_\xi, \alpha|y)$ converges to $p_\infty(\mu, \zeta, \sigma_\xi, \alpha|y)$, the (improper) posterior distribution under $p_\infty(\alpha)$. Thus, by Fatou's lemma, the corresponding conditional distributions also converge, so that, $p_n(\alpha|\mu, \zeta, \sigma_\xi, y) \rightarrow p_\infty(\alpha|\mu, \zeta, \sigma_\xi, y)$. Thus, given $(\mu^{(t+1)}, \zeta^*, \alpha^*)$, Step 3 converges to sampling from the proper distribution

$$\begin{aligned} \alpha^{(t+1)} &\sim \text{N} \left[\frac{\sum_{j=1}^J \zeta_j^* (y_j - \mu^{(t+1)}) / \sigma_j^2}{\sum_{j=1}^J (\zeta_j^*)^2 / \sigma_j^2}, \left(\sum_{j=1}^J (\zeta_j^*)^2 / \sigma_j^2 \right) \right] \\ &= \alpha^* \text{N} \left[\frac{\sum_{j=1}^J \beta_j^* (y_j - \mu^{(t+1)}) / \sigma_j^2}{\sum_{j=1}^J (\beta_j^*)^2 / \sigma_j^2}, \left(\sum_{j=1}^J (\beta_j^*)^2 / \sigma_j^2 \right) \right]. \quad (\text{A.3}) \end{aligned}$$

Notationally, we refer to the normal random variable in (A.3) as α , that is, $\alpha^{(t+1)} = \alpha^* \alpha$.

Finally, in Step 4, we compute $\beta^{(t+1)}$ and $\sigma_\beta^{(t+1)}$ which in the limit simplifies to

$$\beta^{(t+1)} = \alpha^{(t+1)} \zeta^* = \alpha \beta^* \quad \text{and} \quad \sigma_\beta^{(t+1)} = |\alpha^{(t+1)}| \sigma_\zeta^* = |\alpha| \sigma_\beta^*. \quad (\text{A.4})$$

Thus, under the limiting kernel, $\gamma^{(t+1)}$ does not depend on α^* and we do not need to compute α^* , ζ^* , σ_ζ^* , or $\alpha^{(t+1)}$. Thus the iteration consists of sampling steps given in (A.1), (A.2), and (4.1), and computing (A.4). But this is exactly the transition kernel given by the V+PX-SAMPLER. \square

A similar strategy can be used to verify that the S+PX-SAMPLER is the proper Markovian limit of a sequence of proper Markovian transition kernels each with stationary distribution equal to $p(\gamma | y)$. In particular, we use the same sequence of proper prior distributions, $p_n(\alpha)$ to construct the following sequence of transition kernels.

PROPER S+PX-SAMPLER

Step 1: Sample $(\alpha^*, \zeta^*, \sigma_\zeta^*)$ from $p_n(\alpha, \zeta, \sigma_\zeta | \gamma = \gamma^{(t)}, y)$; that is, sample $\alpha^* \sim p_n(\alpha)$ and set $\zeta^* = \beta^{(t)}/\alpha^*$ and $\sigma_\zeta^* = \sigma_\beta^{(t)}/|\alpha^*|$.

Step 2: Sample $\mu^{(t+1)}$ from $p(\mu | \zeta = \zeta^*, \sigma_\zeta = \sigma_\zeta^*, \alpha = \alpha^*, y)$.

Step 3: Sample ζ_j from $p(\zeta_j | \mu = \mu^{(t+1)}, \zeta_{-j}, \sigma_\zeta = \sigma_\zeta^*, \alpha = \alpha^*, y)$ for $j = 1, \dots, J$.

Step 4: Sample σ_ζ from $p(\sigma_\zeta | \mu = \mu^{(t+1)}, \zeta, \alpha = \alpha^*, y)$.

Step 5: Sample $\alpha^{(t+1)}$ from $p_n(\alpha | \mu = \mu^{(t+1)}, \zeta, \sigma_\zeta, y)$.

Step 6: Set $\beta^{(t+1)} = \alpha^{(t+1)} \zeta$ and $\sigma_\beta^{(t+1)} = |\alpha^{(t+1)}| \sigma_\zeta$.

In the limit, we find that $\gamma^{(t+1)}$ does not depend on α_0 , thus the limiting transition kernel is proper and Markovian. Analysis similar to that given for the V+PX-SAMPLER shows that the limiting transition kernel is the kernel described by the S+PX-SAMPLER.

ACKNOWLEDGMENTS

We thank Xiao-Li Meng and several reviewers for helpful discussion and the National Science Foundation (DMS 04-06085) for financial support.

[Received March 2006. Revised July 2007.]

REFERENCES

- Barnard, J., McCulloch, R., and Meng, X. L. (1996), "Modeling Covariance Matrices in Terms of Standard Deviations and Correlations, with Application to Shrinkage," *Statistica Sinica*, 10, 1281–1311.
- Besag, J. (1974), "Spatial Interaction and the Statistical Analysis of Lattice Systems" (with discussion), *Journal of the Royal Statistical Society*, Ser. B, 36, 192–236.

- Besag, J., and Green, P. J. (1993), “Spatial Statistics and Bayesian Computation” (with discussion), *Journal of the Royal Statistical Society*, Ser. B, 55, 25–102.
- Boscardin, W. J. (1996), “Bayesian Analysis for Some Hierarchical Linear Models,” unpublished Ph.D. thesis, Department of Statistics, University of California, Berkeley.
- Boscardin, W. J., and Gelman, A. (1996), “Bayesian Regression with Parametric Models for Heteroscedasticity,” *Advances in Econometrics*, 11A, 87–109.
- Carlin, B. P., and Louis, T. A. (2000), *Bayes and Empirical Bayes Methods for Data Analysis* (2nd ed.), London: Chapman and Hall.
- Daniels, M. J., and Kass, R. E. (1999), “Nonconjugate Bayesian Estimation of Covariance Matrices and its use in Hierarchical Models,” *Journal of the American Statistical Association*, 94, 1254–1263.
- Dempster, A. P., Laird, N. M., and Rubin, D. B. (1977), “Maximum Likelihood from Incomplete Data via the EM Algorithm” (with discussion), *Journal of the Royal Statistical Society*, Ser. B, 39, 1–38.
- Dempster, A. P., Rubin, D. B., and Tsutakawa, R. K. (1981), “Estimation in Covariance Components Models,” *Journal of the American Statistical Association*, 76, 341–353.
- Gelfand, A. E., and Smith, A. F. M. (1990), “Sampling-Based Approaches to Calculating Marginal Densities,” *Journal of the American Statistical Association*, 85, 398–409.
- Gelfand, A. E., Sahu, S. K., and Carlin, B. P. (1995), “Efficient Parameterization for Normal Linear Mixed Models,” *Biometrika*, 82, 479–488.
- Gelman, A. (2004), “Parameterization and Bayesian Modeling,” *Journal of the American Statistical Association*, 99, 537–545.
- (2005), “Analysis of Variance: Why it is More Important than Ever” (with discussion), *The Annals of Statistics*, 33, 1–53.
- (2006), “Prior Distributions for Variance Parameters in Hierarchical Models,” *Bayesian Analysis*, 1, 515–533.
- Gelman, A., Carlin, J. B., Stern, H. S., and Rubin, D. B. (1995), *Bayesian Data Analysis* (1st ed.), London: Chapman and Hall.
- Gelman, A., and Hill, J. (2007), *Data Analysis Using Regression and Multilevel/Hierarchical Models*, New York: Cambridge University Press.
- Gelman, A., and Little, T. C. (1997), “Poststratification into Many Categories using Hierarchical Logistic Regression,” *Survey Methodology*, 23, 127–135.
- Gelman, A., and Rubin, D. B. (1992), “Inference from Iterative Simulation Using Multiple Sequences” (with discussion), *Statistical Science*, 7, 457–511.
- Gilks, W. R., Best, N., and Tan, K. K. C. (1995), “Adaptive Rejection Metropolis Sampling Within Gibbs Sampling,” *Applied Statistics*, 44, 455–472.
- Gilks, W. R., Richardson, S., and Spiegelhalter, D. (eds.) (1996), *Practical Markov Chain Monte Carlo*, London: Chapman and Hall.
- Gilks, W. R., and Roberts, G. O. (1996), “Strategies for Improving MCMC,” in *Practical Markov Chain Monte Carlo*, eds. W. Gilks, S. Richardson, and D. Spiegelhalter, London: Chapman and Hall, pp. 89–114.
- Goldstein, H. (1995), *Multilevel Statistical Models*, London: Edward Arnold.
- Golub, G. H., and van Loan, C. F. (1983), *Matrix Computations*, Baltimore, MD: Johns Hopkins University Press.
- Green, P. J. (1995), “Reversible Jump Markov Chain Monte Carlo Computation and Bayesian Model Determination,” *Biometrika*, 82, 711–732.
- Hills, S. E., and Smith, A. F. M. (1992), “Parametrization Issues in Bayesian Inference” (with discussion), in *Bayesian Statistics 4*, eds. J. M. Bernardo, J. O. Berger, A. P. Dawid, and A. F. M. Smith, New York: Oxford University Press, pp. 227–246.

- Hodges, J. H. (1998), "Some Algebra and Geometry for Hierarchical Models, Applied to Diagnostics" (with discussion), *Journal of the Royal Statistical Society*, Ser. B, 60, 497–536.
- Laird, N. M., and Ware, J. H. (1982), "Random-Effects Models for Longitudinal Data," *Biometrics*, 38, 963–974.
- Lindley, D. V., and Smith, A. F. M. (1972), "Bayes Estimates for the Linear Model," *Journal of the Royal Statistical Society*, Ser. B, 34, 1–41.
- Liu, C. (2003), "Alternating Subspace-Spanning Resampling to Accelerate Markov Chain Monte Carlo Simulation," *Journal of the American Statistical Association*, 98, 110–117.
- Liu, C., Rubin, D. B., and Wu, Y. N. (1998), "Parameter Expansion to Accelerate EM: The PX-EM Algorithm," *Biometrika*, 85, 755–770.
- Liu, J. S. (1994), "The Fraction of Missing Information and Convergence Rate for Data Augmentation," *Computing Science and Statistics*, 26, 490–497.
- Liu, J., and Wu, Y. N. (1999), "Parameter Expansion for Data Augmentation," *Journal of the American Statistical Association*, 94, 1264–1274.
- Longford, N. (1993), *Random Coefficient Models*, Oxford: Clarendon Press.
- MacLehose, R. F., Dunson, D. B., Herring, A., and Hoppin, J. A. (2007), "Bayesian Methods for Highly Correlated Exposure Data," *Epidemiology* 18, 199–207.
- McCullagh, P., and Nelder, J. A. (1989), *Generalized Linear Models* (2nd ed.), London: Chapman and Hall.
- Meng, X. L., and van Dyk, D. (1997), "The EM Algorithm—An Old Folk Song Sung to a Fast New Tune" (with discussion), *Journal of the Royal Statistical Society*, Ser. B, 59, 511–567.
- O'Malley, A. J., and Zaslavsky, A. M. (2005), "Cluster-Level Covariance Analysis for Survey Data with Structured Nonresponse," technical report, Department of Health Care Policy, Harvard Medical School.
- Raftery, A. E. (1996), "Hypothesis Testing and Model Selection via Posterior Simulation," in *Practical Markov Chain Monte Carlo*, eds. W. Gilks, S. Richardson, and D. Spiegelhalter, New York: Chapman and Hall, 163–187.
- Robinson, G. K. (1991), "That BLUP is a Good Thing: The Estimation of Random Effects" (with discussion), *Statistical Science*, 6, 15–51.
- Rosenthal, J. S. (1995), "Minorization Conditions and Convergence Rates for Markov Chain Monte Carlo," *Journal of the American Statistical Association*, 90, 558–566.
- Rubin, D. B. (1981), "Estimation in Parallel Randomized Experiments," *Journal of Educational Statistics*, 6, 377–401.
- Sargent, D. J., Hodges, J. S., and Carlin, B. P. (2000), "Structured Markov Chain Monte Carlo," *Journal of Computational and Graphical Statistics*, 9, 217–234.
- van Dyk, D. A., and Meng, X. L. (2001), "The Art of Data Augmentation" (with discussion), *Journal of Computational and Graphical Statistics*, 10, 1–111.