

Laplace Likelihood and LAD Estimation for Non-invertible MA(1)



Colorado State University
National Tsing-Hua University
U. of California, San Diego

(<http://www.stat.colostate.edu/~rdavis/lectures>)

MA(1) unit root problem

MA(1): (world's simplest time series model!)

$$Y_t = Z_t - \theta Z_{t-1}, \quad \{Z_t\} \sim \text{IID } (0, \sigma^2)$$

Properties:

- $|\theta| < 1 \Rightarrow Z_t = \sum_{j=0}^{\infty} \theta^j Y_{t-j}$ (invertible)
- $|\theta| > 1 \Rightarrow Z_t = -\sum_{j=1}^{\infty} \theta^{-j} Y_{t+j}$ (non-invertible)
- $|\theta| = 1 \Rightarrow Z_t \in \text{sp}\{Y_t, Y_{t-1}, \dots\}$ and $Z_t \in \text{sp}\{Y_{t+1}, Y_{t+2}, \dots\}$
 $\Rightarrow P_{\text{sp}\{Y_s, s \neq 0\}} Y_0 = Y_0$ (perfect interpolation)
- $|\theta| < 1 \Rightarrow \hat{\theta}_{mle}$ is $\text{AN}(\theta, (1 - \theta^2)/n)$

MLE = maximum (Gaussian) likelihood, n = sample size

What if $\theta = 1$?

Why study MA(1) with a unit root?

a) differencing (to remove non-stationarity)

- linear trend model: $X_t = a + bt + Z_t$.

$$Y_t = X_t - X_{t-1} = b + Z_t - Z_{t-1} \sim \text{MA}(1) \text{ with } \theta = 1.$$

- seasonal model: $X_t = s_t + Z_t$, s_t seasonal component w/ period 12.

$$Y_t = X_t - X_{t-12} = Z_t - Z_{t-12} \sim \text{MA}(12) \text{ with } \theta = 1.$$

b) random walk + noise

$$X_t = X_{t-1} + U_t \quad (\text{random walk signal})$$

$$Y_t = X_t + V_t \quad (\text{random walk signal + noise})$$

Then

$$Y_t - Y_{t-1} = U_t + V_t - V_{t-1} \sim \text{MA}(1)$$

with $\theta=1$ if and only if $\text{Var}(U_t) = 0$.

Identifiability and the Gaussian likelihood

Identifiability

- $|\theta| > 1 \Rightarrow Y_t = \varepsilon_t - \theta^{-1} \varepsilon_{t-1}$, where $\{\varepsilon_t\} \sim WN(0, \theta^2 \sigma^2)$.
- $\{\varepsilon_t\}$ is IID if and only if $\{Z_t\}$ is Gaussian (Breidt and Davis '91)
- $\{\varepsilon_t\}$ is a special case of an **All-Pass Model** (Breidt, Davis, Trindade '01, Andrews et al. '05a, '05b)

Gaussian Likelihood

$$L_G(\theta, \sigma^2) = L_G(1/\theta, \theta^2 \sigma^2) \Rightarrow \theta \text{ is only identifiable for } |\theta| \leq 1.$$


Notes:

- i) this implies $L_G(\theta) = L_G(1/\theta)$ for the profile likelihood and $\theta = 1$ is a critical point, $L'_G(1) = 0$.
- ii) a *pile-up effect* ensues, i.e., $P(\hat{\theta} = 1) > 0$ even if $\theta < 1$.

Gaussian likelihood, non-Gaussian data

100 observations from $Y_t = Z_t - \theta_0 Z_{t-1}$, $\{Z_t\} \sim \text{IID}$, Laplace pdf

$$\theta_0 = .8 \quad \theta_0 = 1.0 \quad \theta_0 = 1.25$$



Gaussian MLE for near-unit roots

Idea: build parameter normalization into the likelihood function.

Model: $Y_t = Z_t - (1-\beta/n) Z_{t-1}, t=1,\dots,n.$

$$\beta = n(1-\theta), \theta = 1 - \beta/n, \theta_0 = 1 - \gamma/n$$

Gaussian Likelihood:

$$L_n(\beta) = l_n(1 - \beta/n) - l_n(1), l_n(\) = \text{profile log-like.}$$

Theorem (Davis and Dunsmuir '96): Under $\theta_0 = 1 - \gamma / n$,

$$L_n(\beta) \xrightarrow{d} Z_\gamma(\beta) \text{ on } C[0, \infty).$$

Results:

- $n(1 - \hat{\theta}_{mle}) \rightarrow \hat{\beta}_{mle} = \operatorname{argmax} Z_\gamma(\beta)$
- $n(1 - \hat{\theta}_L) \rightarrow \hat{\beta}_L = \operatorname{arglocalmax} Z_\gamma(\beta)$
- $P(\hat{\theta}_L = 1) \rightarrow P(\hat{\beta}_L = 0) = .6518 \quad \text{if } \gamma = 0.$

Extensions of MLE (Gaussian likelihood)

i) non-zero mean (Chen and Davis '00): same type of limit, except pile-up is more excessive.

$$P(\hat{\theta}_{mle} = 1) \rightarrow .955$$

This makes hypothesis testing easy!

Reject $H_0: \theta = 1$ if $\hat{\theta}_{mle} < 1$ (size of test is .045)

ii) heavy tails (Davis and Mikosch '98): $\{Z_t\}$ symmetric alpha stable (S α S). Then the max Gaussian likelihood estimator has the same normalizing rate, i.e.,

$$n(1 - \hat{\theta}_L) \xrightarrow{d} \hat{\beta}_L$$

$$P(\hat{\theta}_L = 1) \rightarrow P(\hat{\beta}_L = 0)$$

The pile-up decreases with increasing tail heaviness.

Laplace likelihood/LAD estimation

If noise distribution is non-Gaussian, the MA(1) parameter θ is identifiable for all real values.

- Q1. For MLE (non-Gaussian) does one have $1/n$ or $1/n^{1/2}$ asymptotics?
- Q2. Is there a *pile-up* effect?

Look at this problem with *non-Gaussian likelihood*

- Specifically, consider *Laplace likelihood / Least Absolute Deviations* for unit root only (not near-unit root)
- *Preliminary results only!*

Non-Gaussian likelihood – Joint and Exact

Model. $Y_t = Z_t - \theta Z_{t-1}$, $\{Z_t\} \sim \text{IID}$ with median 0 and $EZ^4 < \infty$. **Initial variable.**

$$Z^{init} = \begin{cases} Z_0, & \text{if } |\theta| \leq 1, \\ Z_n - \sum_{t=1}^n Y_t, & \text{otherwise.} \end{cases}$$

Joint density: Let $\mathbf{Y}_n = (Y_1, \dots, Y_n)$, then

$$f(\mathbf{y}_n, z^{init}) = f(z_0, z_1, \dots, z_n) \left(1_{\{|\theta| \leq 1\}} + |\theta|^{-n} 1_{\{|\theta| > 1\}} \right),$$

where the z_t are solved

forward by: $z_t = Y_t + \theta z_{t-1}$, $t = 1, \dots, n$ for $|\theta| \leq 1$ with $z_0 = z^{init}$


backward by: $z_{t-1} = \theta^{-1}(z_t - Y_t)$, $t = n, \dots, 1$ for $|\theta| > 1$ with $z_n = z^{init} + Y_1 + \dots + Y_n$

Note: integrate out z^{init} to get *Exact* likelihood.

$$f(\mathbf{y}_n) = \int_{-\infty}^{\infty} f(\mathbf{y}_n, z^{init}) dz^{init}$$

Laplace likelihood examples

100 observations from $Y_t = Z_t - \theta_0 Z_{t-1}$, $\{Z_t\} \sim \text{IID Laplace pdf}$




Laplace likelihood, Laplace noise


100 observations from $Y_t = Z_t - \theta_0 Z_{t-1}$, $\{Z_t\} \sim \text{IID Laplace pdf}$

$$\theta_0 = .8 \quad \theta_0 = 1.0 \quad \theta_0 = 1.25$$

Exact likelihood



Joint likelihood at $z_{\max}(\theta)$



Laplace likelihood-LAD estimation

(Joint) Laplace log-likelihood. ($\sigma = E|Z_0|$ is a scale parameter)

$$L(\theta, z^{init}, \sigma) = -(n+1)\log 2\sigma - \sigma^{-1} \sum_{t=0}^n |z_t| - n(\log |\theta|)1_{\{|\theta|>1\}}$$

Maximizing wrt σ , we obtain

$$\hat{\sigma} = \sum_{t=0}^n |z_t| / (n+1)$$

so that maximizing L is equivalent to minimizing

$$l_n(\theta, z^{init}) = \begin{cases} \sum_{t=0}^n |z_t|, & \text{if } |\theta| \leq 1, \\ \sum_{t=0}^n |z_t| |\theta|, & \text{otherwise.} \end{cases}$$

Joint Laplace likelihood — limit results

Result 1. Under the parameterizations,

$$\theta = 1 + \beta/n \quad \text{and} \quad z^{\text{init}} = Z_0 + \alpha\sigma/n^{1/2},$$

we have

$$U_n(\beta, \alpha) = \sigma^{-1}(l_n(\theta, z^{\text{init}}) - l_n(1, Z_0)) \rightarrow_d U(\beta, \alpha)$$

on $C(\mathbb{R}^2)$, where

$$\begin{aligned} U(\beta, \alpha) &= \int_0^1 \left(\beta \int_0^s e^{\beta(s-t)} dS(t) + \alpha e^{\beta s} \right) dW(s) \\ &\quad + f(0) \int_0^1 \left(\beta \int_0^s e^{\beta(s-t)} dS(t) + \alpha e^{\beta s} \right)^2 ds \end{aligned}$$

for $\beta \leq 0$, and

$$\begin{aligned} U(\beta, \alpha) &= \int_0^1 \left(-\beta \int_{s+}^1 e^{\beta(s-t)} dS(t) + \alpha e^{-\beta(1-s)} \right) dW(s) \\ &\quad + f(0) \int_0^1 \left(\beta \int_s^1 e^{\beta(s-t)} dS(t) + \alpha e^{-\beta(1-s)} \right)^2 ds \end{aligned}$$

for $\beta > 0$.

Joint Laplace likelihood — limit results

The limits contain correlated Brownian Motions $S(t)$ and $W(t)$, obtained as the limits of the partial sum processes

$$S_n(t) = \frac{1}{\sigma\sqrt{n}} \sum_{i=0}^{[nt]} Z_i \rightarrow_d S(t), \quad W_n(t) = \frac{1}{\sigma\sqrt{n}} \sum_{i=0}^{[nt]} \text{sign}(Z_i) \rightarrow_d W(t).$$

From the limit,

$$U_n(\beta, \alpha) \rightarrow_d U(\beta, \alpha),$$

it suggests (from the continuous mapping theorem?) that

$$\text{limit(optimum(criterion))} = \text{optimum(limit(criterion))}.$$

So for the Local optimizer of the Joint likelihood

$$\left(n \left(\hat{\theta}_{LJ} - 1 \right) \sqrt{n} \sigma^{-1} \left(\hat{z}_{LJ}^{\text{init}} - Z_0 \right) \right) \rightarrow_d \left(\hat{\beta}_{LJ}, \hat{\alpha}_{LJ} \right)$$

where

$$(\hat{\beta}_{LJ}, \hat{\alpha}_{LJ}) = \arg(\text{local}) \min U(\beta, \alpha).$$

Joint Laplace likelihood — limit results

Might expect a similar result to hold for the **Global optimizer** of the **Joint likelihood**

$$\left(n \left(\hat{\theta}_{\text{GJ}} - 1 \right), \sqrt{n} \sigma^{-1} \left(\hat{z}_{\text{GJ}}^{\text{init}} - Z_0 \right) \right) \xrightarrow{d} \left(\hat{\beta}_{\text{GJ}}, \hat{\alpha}_{\text{GJ}} \right)$$

where

$$(\hat{\beta}_{\text{GJ}}, \hat{\alpha}_{\text{GJ}}) = \operatorname{argmin} U(\beta, \alpha).$$

Exact Laplace likelihood — limit results

Exact Laplace Likelihood:

$$L_n(\theta, \sigma) = \int_{-\infty}^{\infty} f(\mathbf{y}_n, z^{init}) dz^{init}$$

Result 2. For the **Global** optimizer of the **Exact** likelihood,

$$n(\hat{\theta}_{GE} - 1) \xrightarrow{d} \hat{\beta}_{GE},$$

where

$$\hat{\beta}_{GE} = \arg \min U^*(\beta),$$

and $U^*(\beta)$ is a stochastic process defined in terms of $S(t)$ and $W(t)$.

In addition, for the **Local** optimizer of the **Exact** likelihood

$$n(\hat{\theta}_{LE} - 1) \xrightarrow{d} \hat{\beta}_{LE}, \quad \hat{\beta}_{LE} = \arg (\text{local}) \min U^*(\beta).$$

Simulating from the limit process

Step 1. Simulate two indep sequences (W_1, \dots, W_m) and (V_1, \dots, V_m) of iid $N(0,1)$ random variables with $m=100000$.

Step 2. Form $W(t)$ and $V(t)$ by the partial sum processes,

$$W(t) = \sum_{j=1}^{[100000t]} W_j / \sqrt{100000} \quad \text{and} \quad V(t) = \sum_{j=1}^{[100000t]} V_j / \sqrt{100000}.$$

Step 3. Set $S(t) = W(t) + c_1 V(t)$, where

$$c_1 = \sqrt{\text{Var}(Z_t)/E^2 |Z_0| - 1}.$$

Limit process depends only on c_1 and $f(0)$.

Step 4. Compute $U(\beta,\alpha)$ and $U^*(\beta)$ from the definition.


Step 5. Determine the respective Local and Global minimizers of Joint limit $U(\beta,\alpha)$ and Exact limit $U^*(\beta)$ numerically.

Simulated realizations of the limit processes

Simulate Joint and Exact limit processes, $U(\beta, \alpha(\beta))$, $U^*(\beta)$.


- Simulate realization of each limit process, joint and exact
- Compute local and global optima
- Repeat...
- Build up limit distribution functions

$t(5)$ pdf




Limit cdf

red graph = Laplace pdf for Z_t



blue graph = Gaussian pdf for Z_t



Local results

Laplace noise

$\theta = 1, \sigma = 1$

1000 reps

n		Exact	Joint	
		$\hat{\theta}_{LE}$	$\hat{\theta}_{LJ}$	$\hat{\sigma}$
$n = 20$	bias	-.0057	-.0033	-.0208
	s.d.	.1438	.0656	.2430
	rmse	.1439	.0657	.2438
	asympt	.1207	.0526	.2236
$n = 50$	bias	.0000	.0004	.0293
	s.d.	.0574	.0208	.1511
	rmse	.0574	.0208	.1539
	asympt	.0483	.0211	.1414
$n = 100$	bias	.0005	-.0003	-.0025
	s.d.	.0303	.0107	.1000
	rmse	.0303	.0107	.1000
	asympt	.0241	.0105	.1000
$n = 200$	bias	.0005	.0000	-.0016
	s.d.	.0140	.0058	.0718
	rmse	.0141	.0058	.0718
	asympt	.0121	.0053	.0707

Simulation results: Global Exact and Global Joint

Global Exact = MLE

Global Joint = maximize over θ and z_{init}

Laplace noise

$\theta = 1, \sigma = 1$

1000 reps

Note:


- Local dominates Global
- Joint dominates Exact
(rmse is half the size)

n		Exact $\hat{\theta}_{GE}$	Joint $\hat{\theta}_{GJ}$	Local $\hat{\theta}_{LJ}$
$n = 20$	bias	-.047	-.050	-.003
	rmse	.224	.213	.144
$n = 50$	bias	-.013	.002	.000
	rmse	.096	.078	.057
$n = 100$	bias	.003	-.003	.000
	rmse	.051	.034	.011
$n = 200$	bias	.000	.000	.000
	rmse	.028	.014	.006

Analysis of pile-up probabilities

Look back at realizations of the limit processes, $U(\beta, \alpha(\beta))$, $U^*(\beta)$.

- When is there a local optimum at $\theta = 1$?
- Check derivatives
- Negative derivative from the left
- Positive derivative from the right
- Local optimum at $\theta = 1$



Pile-up probabilities (Joint)

Result 3. (Local Joint Laplace likelihood)

$$P(\hat{\theta}_{LJ} = 1) \rightarrow P(0 < Y < 1),$$

where

$$Y = \int_0^1 S(s)dW(s) - W(1) \int_0^1 S(s)ds + \frac{W(1)}{2f(0)} \left(\int_0^1 W(s)ds - W(1)/2 \right)$$

Idea: look at derivatives

$$\begin{aligned} P(\hat{\theta}_{lm} = 1) &= P\left(\lim_{\beta \uparrow 0} \frac{\partial}{\partial \beta} U_n(\beta, \hat{\alpha}(\beta)) < 0 \text{ and } \lim_{\beta \downarrow 0} \frac{\partial}{\partial \beta} U_n(\beta, \hat{\alpha}(\beta)) > 0\right) \\ &\rightarrow P\left(\lim_{\beta \uparrow 0} \frac{\partial}{\partial \beta} U(\beta, \hat{\alpha}(\beta)) < 0 \text{ and } \lim_{\beta \downarrow 0} \frac{\partial}{\partial \beta} U(\beta, \hat{\alpha}(\beta)) > 0\right) \end{aligned}$$

Now,

$$\lim_{\beta \downarrow 0} \frac{\partial}{\partial \beta} U(\beta, \hat{\alpha}(\beta)) = Y$$

$$\lim_{\beta \uparrow 0} \frac{\partial}{\partial \beta} U(\beta, \hat{\alpha}(\beta)) = Y - 1 \quad \text{and the result follows.}$$

Pile-up probabilities (Exact)

Result 4. (Local Exact Laplace likelihood)

$$P(\hat{\theta}_{LE} = 1) \rightarrow P\left[\frac{1}{2} < Y < 1 - \frac{1}{2}\right] = 0$$

The *pile-up probability* is always *zero* for the Local Exact, and always positive for the Local Joint (see Result 3).

Remark. (Laplace pile-up)

If Z_t has a Laplace density $f(z) = \frac{1}{2\sigma} e^{-|z|/\sigma}$, then

$$Y = \int_0^1 [W(1)s - W(s)] dV(s) + \frac{1}{2}.$$

where $W(s)$ and $V(s)$ are independent standard Brownian motions.

Laplace pile-up probabilities (cont)

It follows that Local Joint has pile-up probability

$$\begin{aligned} P(\hat{\theta}_{LJ} = 1) &\rightarrow P(0 < Y < 1) \\ &= P(0 < \int_0^1 [W(1)s - W(s)] dV(s) + .5 < 1) \\ &= E \left[P(-.5 < \int_0^1 [W(1)s - W(s)] dV(s) < .5 | W(t), t \in [0,1]) \right] \\ &= E \left[2\Phi \left(.5 \left\{ \int_0^1 [W(1)s - W(s)]^2 ds \right\}^{-1/2} \right) - 1 \right] \\ &\approx 0.820 \end{aligned}$$

But *no* pile-up probability for Local Exact:

Remark: if Local *does not* pile up, Global does not pile up

if Local *does* pile up, Global probably does as well

Simulation results – pile-up probabilities

Pile-up probabilities for **Local Joint**: $P(\hat{\theta}_{LJ} = 1)$

<i>n</i>	Gau	Lap	Unif	t(5)
20	.827	.796	.831	.796
50	.859	.806	.864	.823
100	.873	.819	.864	.817
200	.844	.819	.843	.831
500	.855	.809	.841	.846
∞	.858	.820	.836	.827

(No pile-up probabilities for **Local Exact**.)