

Application of the Innovations Algorithm to Nonlinear State-Space Models

Richard A. Davis
Colorado State University

(<http://www.stat.colostate.edu/~rdavis/lectures>)

Joint work with:

William Dunsmuir, University of New South Wales

Gabriel Rodriguez-Yam, Colorado State University

Ying Wang, Dept of Public Health, W. Virginia



Two CSU Statistics Department Projects

STARMAP

Space-Time Aquatic Resources Modeling and Analysis Program (funded by the EPA-STAR program)

PRIMES

Program for Interdisciplinary Mathematics, Ecology, and Statistics (recommended for funding by NSF IGERT)

- Student fellowships
- Postdoctoral fellowships
- Short-term visitors
- Workshops/seminars and more.

- Generalized state-space models
 - Observation driven
 - Parameter driven
- Innovations algorithm (recursive one-step ahead prediction algorithm)
 - Applications
 - Gaussian likelihood calculations
 - simulation
 - generalized least squares estimation
- Time series of counts
 - Examples (asthma data, polio data)
 - Generalized linear models (GLM)
 - Estimating equations (Zeger)
 - MCEM (Chan and Ledolter)
 - Importance sampling
 - Durbin and Koopman
 - Approximation to the likelihood (Davis, Dunsmuir, and Wang)
 - Simulation results
- Examples

Generalized State-Space Models (parameter driven)

Observations: $\mathbf{y}^{(t)} = (y_1, \dots, y_t)$

States: $\boldsymbol{\alpha}^{(t)} = (\alpha_1, \dots, \alpha_t)$

Observation equation:

$$p(y_t | \alpha_t) := p(y_t | \alpha_t, \boldsymbol{\alpha}^{(t-1)}, \mathbf{y}^{(t-1)})$$

State equation:

$$p(\alpha_{t+1} | \alpha_t) := p(\alpha_{t+1} | \alpha_t, \boldsymbol{\alpha}^{(t-1)}, \mathbf{y}^{(t)})$$

Joint density:

$$\begin{aligned} & p(y_1, \dots, y_n, \alpha_1, \dots, \alpha_n) \\ &= p(y_n | \alpha_n, \boldsymbol{\alpha}^{(n-1)}, \mathbf{y}^{(n-1)}) p(\alpha_n, \boldsymbol{\alpha}^{(n-1)}, \mathbf{y}^{(n-1)}) \\ &= p(y_n | \alpha_n) p(\alpha_n | \boldsymbol{\alpha}^{(n-1)}, \mathbf{y}^{(n-1)}) p(\boldsymbol{\alpha}^{(n-1)}, \mathbf{y}^{(n-1)}) \\ &= \dots \\ &= \left(\prod_{j=1}^n p(y_j | \alpha_j) \right) \left(\prod_{j=2}^n p(\alpha_j | \alpha_{j-1}) \right) p(\alpha_1) \end{aligned}$$

Parameter driven (cont)

Conditional independence:

$$p(y_1, \dots, y_n | \alpha_1, \dots, \alpha_n) = \prod_{j=1}^n p(y_j | \alpha_j)$$

Filtering or posterior density:

$$p(\alpha_t | \mathbf{y}^{(t)}) = p(y_t | \alpha_t) p(\alpha_t | \mathbf{y}^{(t-1)}) / p(y_t | \mathbf{y}^{(t-1)})$$

Predictive densities:

$$p(\alpha_{t+1} | \mathbf{y}^{(t)}) = \int p(\alpha_t | \mathbf{y}^{(t)}) p(\alpha_{t+1} | \alpha_t) d\mu(\alpha_t)$$

$$p(y_{t+1} | \mathbf{y}^{(t)}) = \int p(y_{t+1} | \alpha_{t+1}) p(\alpha_{t+1} | \mathbf{y}^{(t)}) d\mu(\alpha_{t+1})$$

Examples of parameter driven models

Poisson model for time series of counts

Observation equation:

$$p(y_t | \alpha_t) = \frac{e^{\alpha_t y_t} e^{-e^{\alpha_t}}}{y_t!}, \quad y_t = 0, 1, \dots,$$

State equation: State variables follow a regression model with Gaussian AR(1) noise

$$\alpha_t = \beta^T \mathbf{x}_t + W_t, \quad W_t = \phi W_{t-1} + Z_t, \quad \{Z_t\} \sim \text{WN}(0, \sigma^2)$$

The resulting transition density of the state variables is

$$p(\alpha_{t+1} | \alpha_t) = n(\alpha_{t+1}; \beta^T \mathbf{x}_{t+1} + \phi (\alpha_t - \beta^T \mathbf{x}_t), \sigma^2)$$

Remark: The case $\sigma^2 = 0$ corresponds to a log-linear model with Poisson noise.

Examples of parameter driven models

A stochastic volatility model for financial data (Taylor `86):

Model:

$$Y_t = \sigma_t Z_t, \{Z_t\} \sim \text{IID } N(0,1)$$

$$\alpha_t = \phi \alpha_{t-1} + W_t, \{W_t\} \sim \text{IID } N(0, \sigma^2),$$

where $\alpha_t = \log \sigma_t$.

The resulting observation and state transition densities are

$$p(y_t | \alpha_t) = n(y_t; 0, \exp(2\alpha_t))$$

$$p(\alpha_{t+1} | \alpha_t) = n(\alpha_{t+1}; \phi \alpha_t, \sigma^2)$$

Properties:

- Martingale difference sequence.
- Stationary.
- Strongly mixing at a geometric rate.

The Innovations Algorithm

Innovations Algorithm (Brockwell and Davis '87): $\{X_t\}$ is a zero-mean time series with ACVF $\kappa(i,j)$, then

$$\hat{X}_{t+1} = P_{\text{sp}\{1, X_1, \dots, X_t\}} X_{t+1} = \theta_{t1} (X_t - \hat{X}_t) + \dots + \theta_{tt} (X_1 - \hat{X}_1)$$

The coefficients $\theta_{t1}, \dots, \theta_{tt}$ and prediction errors v_{t-1} can be computed recursively from the equations,

$$v_0 = \kappa(1,1)$$

$$\theta_{t,t-k} = \left[\kappa(t+1, k+1) - \sum_{j=0}^{k-1} \theta_{k,k-j} \theta_{t,t-j} v_j \right] v_{k-1}^{-1}, \quad k = 0, \dots, t-1,$$

and

$$v_t = \kappa(t+1, t+1) - \sum_{j=0}^{t-1} \theta_{t,t-j}^2 v_j.$$

The Innovations Algorithm(cont)

Remarks:

- Innovations algorithm expresses one-step predictor in terms of previous *innovations*, $X_1 - \hat{X}_1, \dots, X_t - \hat{X}_t$, that are uncorrelated.

- If $\{X_t\}$ is an MA(q) process

$$X_{t+1} = Z_{t+1} + \theta_1 Z_t + \dots + \theta_q Z_{t-q}, \quad \{Z_t\} \sim WN(0, \sigma^2)$$

then $(\theta_{t1}, \dots, \theta_{tt}) = (\theta_{t1}, \dots, \theta_{tq}, 0, \dots, 0)$ for all t .

- Innovations algorithm is well adapted for ARMA(p,q) models—only need to apply to MA(q) piece (see B&D `96).

The Innovations Algorithm—Applications

Likelihood calculation:

Using the IA representation,

$$\hat{X}_t = \theta_{t-1,1}(X_{t-1} - \hat{X}_{t-1}) + \cdots + \theta_{t-1,t-1}(X_1 - \hat{X}_1)$$

we have

$$\begin{bmatrix} X_1 \\ X_2 \\ X_3 \\ \vdots \\ X_n \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ \theta_{1,1} & 1 & 0 & \cdots & 0 \\ \theta_{2,2} & \theta_{2,1} & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \theta_{n-1,n-1} & \theta_{n-1,n-2} & \theta_{n-1,n-3} & \cdots & 1 \end{bmatrix} \begin{bmatrix} X_1 - \hat{X}_1 \\ X_2 - \hat{X}_2 \\ X_3 - \hat{X}_3 \\ \vdots \\ X_n - \hat{X}_n \end{bmatrix}$$

$$\mathbf{X}_n = \mathbf{C}_n (\mathbf{X}_n - \hat{\mathbf{X}}_n)$$

By taking covariances of both sides it follows that

$$\Gamma_n = E(\mathbf{X}_n \mathbf{X}_n') = \mathbf{C}_n \mathbf{D}_n \mathbf{C}_n', \quad \mathbf{D}_n = \text{diag}(v_0, \dots, v_{n-1})$$

The Innovations Algorithm—Applications

Quadratic form:

$$\begin{aligned}\mathbf{X}'_n \Gamma_n^{-1} \mathbf{X}_n &= (\mathbf{X}_n - \hat{\mathbf{X}}_n)' C'_n (C_n^{-1} D_n^{-1} C_n) C_n (\mathbf{X}_n - \hat{\mathbf{X}}_n) \\ &= (\mathbf{X}_n - \hat{\mathbf{X}}_n)' D_n^{-1} (\mathbf{X}_n - \hat{\mathbf{X}}_n) \\ &= \sum_{t=1}^n (X_t - \hat{X}_t)^2 / v_{t-1}\end{aligned}$$

Determinant:

$$\det(\Gamma_n) = \det(C_n D_n C'_n) = v_0 \cdots v_{n-1}$$

Gaussian likelihood:

$$L(\Gamma_n) = (2\pi)^{-n/2} (v_0 \cdots v_{n-1})^{-1/2} \exp\{-1/2 \sum_{t=1}^n (X_t - \hat{X}_t)^2 / v_{t-1}\}$$

Simulation: If $\{Z_t\} \sim \text{iid } N(0,1)$, put $X_t = v_{t-1}^{-1/2} Z_t + \theta_{t-1,1} v_{t-2}^{-1/2} Z_{t-1} + \cdots + \theta_{t-1,t-1} v_0^{-1/2} Z_1$.

Then $\mathbf{X}_n = (X_1, \dots, X_n)' = C'_n D_n^{-1/2} \mathbf{Z}_n$

has covariance matrix Γ_n .

Time Series of Counts—Notation and Setup

Count data: Y_1, \dots, Y_n

Regression (explanatory) variable: \mathbf{x}_t

Model: Distribution of the Y_t given \mathbf{x}_t and a stochastic process α_t are indep

Poisson distributed with mean





$$\mu_t = \exp(\mathbf{x}_t^T \boldsymbol{\beta} + \alpha_t).$$

The distribution of the stochastic process α_t may depend on a vector of parameters $\boldsymbol{\gamma}$.


Note: $\alpha_t = 0$ corresponds to standard Poisson regression model.

Primary objective: Inference about $\boldsymbol{\beta}$.

Example: Daily Asthma Presentations (1990:1993)



Polio Data With Estimated Regression Function



Parameter-Driven Model for the Mean Function μ_t

Parameter-driven specification: (Assume $Y_t | \mu_t$ is Poisson(μ_t))

$$\log \mu_t = \mathbf{x}_t^T \boldsymbol{\beta} + \alpha_t ,$$

where $\{\alpha_t\}$ is a stationary Gaussian process.

e.g. (AR(1) process)

$$(\alpha_t + \sigma^2/2) = \phi(\alpha_{t-1} + \sigma^2/2) + \varepsilon_t , \quad \{\varepsilon_t\} \sim \text{IID } N(0, \sigma^2(1-\phi^2)).$$

Advantages of this model specification:

- properties of model (ergodicity and mixing) easy to derive.
- interpretability of regression parameters

$$E(Y_t) = \exp(\mathbf{x}_t^T \boldsymbol{\beta}) E \exp(\alpha_t) = \exp(\mathbf{x}_t^T \boldsymbol{\beta}) , \quad \text{if } E \exp(\alpha_t) = 1.$$

Disadvantages:

- estimation is difficult-likelihood function not easily calculated (MCEM, importance sampling, estimating eqns).
- model building can be laborious

Remark: See Davis, Dunsmuir, and Wang (1999) for testing of the existence of a latent process and estimating its ACF.

Estimation Methods — Importance Sampling (Durbin and Koopman)

Model:

$$Y_t | \alpha_t, \mathbf{x}_t \sim \text{Pois}(\exp(\mathbf{x}_t^T \boldsymbol{\beta} + \alpha_t))$$

$$\alpha_t = \phi \alpha_{t-1} + \varepsilon_t, \quad \{\varepsilon_t\} \sim \text{IID } N(0, \sigma^2)$$

Relative Likelihood: Let $\boldsymbol{\psi} = (\boldsymbol{\beta}, \phi, \sigma^2)$ and suppose $g(\mathbf{y}_n, \boldsymbol{\alpha}_n; \boldsymbol{\psi}_0)$ is an approximating joint density for $\mathbf{Y}_n = (Y_1, \dots, Y_n)'$ and $\boldsymbol{\alpha}_n = (\alpha_1, \dots, \alpha_n)'$.

$$\begin{aligned} L(\boldsymbol{\psi}) &= \int p(\mathbf{y}_n | \boldsymbol{\alpha}_n) p(\boldsymbol{\alpha}_n) d\boldsymbol{\alpha}_n \\ &= \int \frac{p(\mathbf{y}_n | \boldsymbol{\alpha}_n) p(\boldsymbol{\alpha}_n)}{g(\mathbf{y}_n, \boldsymbol{\alpha}_n; \boldsymbol{\psi}_0)} g(\mathbf{y}_n, \boldsymbol{\alpha}_n; \boldsymbol{\psi}_0) d\boldsymbol{\alpha}_n \\ &= \int \frac{p(\mathbf{y}_n | \boldsymbol{\alpha}_n) p(\boldsymbol{\alpha}_n)}{g(\mathbf{y}_n, \boldsymbol{\alpha}_n; \boldsymbol{\psi}_0)} g(\boldsymbol{\alpha}_n | \mathbf{y}_n; \boldsymbol{\psi}_0) g(\mathbf{y}_n; \boldsymbol{\psi}_0) d\boldsymbol{\alpha}_n \\ \frac{L(\boldsymbol{\psi})}{L_g(\boldsymbol{\psi}_0)} &= \int \frac{p(\mathbf{y}_n | \boldsymbol{\alpha}_n) p(\boldsymbol{\alpha}_n)}{g(\mathbf{y}_n, \boldsymbol{\alpha}_n; \boldsymbol{\psi}_0)} g(\boldsymbol{\alpha}_n | \mathbf{y}_n; \boldsymbol{\psi}_0) d\boldsymbol{\alpha}_n \end{aligned}$$

Importance Sampling (cont)

$$\begin{aligned}\frac{L(\psi)}{L_g(\psi_0)} &= \int \frac{p(\mathbf{y}_n | \boldsymbol{\alpha}_n) p(\boldsymbol{\alpha}_n)}{g(\mathbf{y}_n, \boldsymbol{\alpha}_n; \psi_0)} g(\boldsymbol{\alpha}_n | \mathbf{y}_n; \psi_0) d\boldsymbol{\alpha}_n \\ &= E_g \left[\frac{p(\mathbf{y}_n | \boldsymbol{\alpha}_n) p(\boldsymbol{\alpha}_n)}{g(\mathbf{y}_n, \boldsymbol{\alpha}_n; \psi_0)} \mid \mathbf{y}_n; \psi_0 \right] \\ &\sim \frac{1}{N} \sum_{j=1}^N \frac{p(\mathbf{y}_n | \boldsymbol{\alpha}_n^{(j)}) p(\boldsymbol{\alpha}_n^{(j)})}{g(\mathbf{y}_n, \boldsymbol{\alpha}_n^{(j)}; \psi_0)},\end{aligned}$$

where $\{\boldsymbol{\alpha}_n^{(j)}; j = 1, \dots, N\} \sim \text{iid } g(\boldsymbol{\alpha}_n | \mathbf{y}_n; \psi_0)$.





Notes:

- This is a “one-sample” approximation to the relative likelihood. That is, for one realization of the $\boldsymbol{\alpha}$'s, we have, in principle, an approximation to the whole likelihood function.
- Approximation is only good in a neighborhood of ψ_0 . Geyer suggests maximizing ratio wrt ψ and iterate replacing ψ_0 with $\hat{\psi}$.










Importance Sampling — example

Simulation example: $Y_t | \alpha_t \sim \text{Pois}(\exp(.7 + \alpha_t))$,

$$\alpha_t = .5 \alpha_{t-1} + \varepsilon_t, \quad \{\varepsilon_t\} \sim \text{IID } N(0, .3), \quad n = 200, \quad N = 1000$$



Simulation example: $Y_t | \alpha_t \sim \text{Pois}(\exp(.7 + \alpha_t))$, $\phi = .5$, $\sigma^2 = .3$, $n = 200$, $N = 1000$



Importance Sampling (cont)

Choice of *importance density* g :

Durbin and Koopman suggest a linear state-space approximating model

$$Y_t = \mu_t + \mathbf{x}_t^T \boldsymbol{\beta} + \alpha_t + Z_t, \quad Z_t \sim N(0, H_t),$$

with

$$\begin{aligned} \mu_t &= y_t - \hat{\alpha}_t - \mathbf{x}_t' y_t e^{-(\hat{\alpha}_t + \mathbf{x}_t' \boldsymbol{\beta})} + 1, \\ H_t &= e^{-(\hat{\alpha}_t + \mathbf{x}_t' \boldsymbol{\beta})}, \end{aligned}$$

where the $\hat{\alpha}_t = E_g(\alpha_t | \mathbf{y}_n)$ are calculated recursively under the approximating model until convergence.

With this choice of approximating model, it turns out that

$$g(\boldsymbol{\alpha}_n | \mathbf{y}_n; \Psi_0) \sim N(\Gamma_n^{-1} \tilde{\mathbf{y}}_n, \Gamma_n^{-1}),$$

where

$$\begin{aligned} \tilde{\mathbf{y}}_n &= \mathbf{y}_n - e^{\mathbf{X}\boldsymbol{\beta} + \hat{\boldsymbol{\alpha}}_n} + e^{\mathbf{X}\boldsymbol{\beta} + \hat{\boldsymbol{\alpha}}_n} \hat{\boldsymbol{\alpha}}_n, \\ \Gamma_n &= \text{diag}(e^{\mathbf{X}\boldsymbol{\beta} + \hat{\boldsymbol{\alpha}}_n}) + (E(\boldsymbol{\alpha}_n \boldsymbol{\alpha}_n'))^{-1}. \end{aligned}$$

Importance Sampling (cont)

Components required in the calculation.

- $g(\mathbf{y}_n, \boldsymbol{\alpha}_n)$
 - ◆ $\tilde{\mathbf{y}}_n' \Gamma_n^{-1} \tilde{\mathbf{y}}_n$
 - ◆ $\det(\Gamma_n)$
- simulate from $N(\Gamma_n^{-1} \tilde{\mathbf{y}}_n, \Gamma_n^{-1})$
 - ◆ compute $\Gamma_n^{-1} \tilde{\mathbf{y}}_n$
 - ◆ simulate from $N(\mathbf{0}, \Gamma_n^{-1})$

Importance Sampling (cont)

Details.

$$(E(\mathbf{a}_n \mathbf{a}'_n))^{-1} = \sigma^{-2} \begin{pmatrix} 1 & -\phi & 0 & \cdots & 0 \\ -\phi & 1+\phi^2 & -\phi & \cdots & 0 \\ 0 & -\phi & 1+\phi^2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1+\phi^2 \end{pmatrix}$$

$$\Gamma_n = \text{diag}(e^{\hat{\alpha} + X\beta}) + \sigma^{-2} \begin{pmatrix} 1 & -\phi & 0 & \cdots & 0 \\ -\phi & 1+\phi^2 & -\phi & \cdots & 0 \\ 0 & -\phi & 1+\phi^2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1+\phi^2 \end{pmatrix}.$$

This is the covariance function of a 1-dependent sequence, so that $\Gamma_n = C_n D_n C'_n$,

where

$$C_n = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ \theta_{1,1} & 1 & 0 & \cdots & 0 \\ 0 & \theta_{2,1} & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}.$$

Importance Sampling (cont)

It follows that

$$\tilde{\mathbf{y}}_n' \Gamma_n^{-1} \tilde{\mathbf{y}}_n = \sum_{t=1}^n (\tilde{y}_t - \hat{y}_t)^2 / v_{t-1}$$

and

$$\begin{aligned} \Gamma_n^{-1} \tilde{\mathbf{y}}_n &= C_n'^{-1} D_n^{-1} C_n^{-1} C_n (\tilde{\mathbf{y}}_n - \hat{\mathbf{y}}_n) \\ &= C_n'^{-1} (D_n^{-1} (\tilde{\mathbf{y}}_n - \hat{\mathbf{y}}_n)) \end{aligned}$$

which can be solved for the vector $\Gamma_n^{-1} \tilde{\mathbf{y}}_n$ via the recursion

$$C_n' \Gamma_n^{-1} \tilde{\mathbf{y}}_n = D_n^{-1} (\tilde{\mathbf{y}}_n - \hat{\mathbf{y}}_n).$$

All of these calculations can be carried out quickly using the *innovations algorithm*.


To simulate from $N(\mathbf{0}, \Gamma_n^{-1})$ note that

$$\mathbf{U}_n = C_n'^{-1} D_n^{-1} \mathbf{Z}_n,$$

where $\mathbf{Z}_n \sim N(0,1)$, has covariance matrix Γ_n^{-1} .

Importance Sampling — example

Simulation example: $\beta = .7$, $\phi = .5$, $\sigma^2 = .3$, $n = 200$, $N = 1000$, 50 realizations plotted



Estimation Methods — Approximation to the likelihood

Joint density function:

$$p(\mathbf{y}_n, \mathbf{a}_n) \propto \frac{\det(G)^{1/2}}{\prod_{t=1}^n y_t!} \exp\{-(\mathbf{y}_n^T (\mathbf{a}_n + \mathbf{X}\boldsymbol{\beta}) - e^{\mathbf{1}^T (\mathbf{a}_n + \mathbf{X}\boldsymbol{\beta})} - \mathbf{a}_n^T G_n \mathbf{a}_n / 2)\},$$

where $G_n^{-1} = E(\mathbf{a}_n^T \mathbf{a}_n)$.

Conditional density function:

$$p(\mathbf{a}_n | \mathbf{y}_n) \propto \exp\{-\mathbf{y}_n^T \mathbf{a}_n - e^{\mathbf{1}^T (\mathbf{a}_n + \mathbf{X}\boldsymbol{\beta})} - \mathbf{a}_n^T G_n \mathbf{a}_n / 2\},$$

which, by expanding the term, $e^{\mathbf{1}^T (\mathbf{a}_n + \mathbf{X}\boldsymbol{\beta})}$ in a neighborhood of \mathbf{a}_n^* , and ignoring third-order + terms yields the approximation

$$p_a(\mathbf{a}_n | \mathbf{y}_n) \propto \exp\{-(\mathbf{y}_n^T (\mathbf{a}_n + \mathbf{X}\boldsymbol{\beta}) - e^{\mathbf{1}^T (\mathbf{a}_n^* + \mathbf{X}\boldsymbol{\beta})} + (\mathbf{a}_n - \mathbf{a}_n^*)^T e^{\mathbf{a}_n^* + \mathbf{X}\boldsymbol{\beta}} + \frac{1}{2} (\mathbf{a}_n - \mathbf{a}_n^*)^T \text{diag}(e^{\mathbf{a}_n^* + \mathbf{X}\boldsymbol{\beta}}) (\mathbf{a}_n - \mathbf{a}_n^*) - \mathbf{a}_n^T G_n \mathbf{a}_n / 2\}.$$

Estimation Methods — Approximation to the likelihood

After simplification, we find

$$\begin{aligned} p_a(\boldsymbol{\alpha}_n | \mathbf{y}_n) &\propto \exp\{-(\mathbf{y}_n^T (\boldsymbol{\alpha}_n + \mathbf{X}\boldsymbol{\beta}) - e^{1^T (\boldsymbol{\alpha}_n^* + \mathbf{X}\boldsymbol{\beta})} + (\boldsymbol{\alpha}_n - \boldsymbol{\alpha}_n^*)^T e^{\boldsymbol{\alpha}_n^* + \mathbf{X}\boldsymbol{\beta}} \\ &\quad + \frac{1}{2} (\boldsymbol{\alpha}_n - \boldsymbol{\alpha}_n^*)^T \text{diag}(e^{\boldsymbol{\alpha}_n^* + \mathbf{X}\boldsymbol{\beta}}) (\boldsymbol{\alpha}_n - \boldsymbol{\alpha}_n^*) - \boldsymbol{\alpha}_n^T G_n \boldsymbol{\alpha}_n / 2\}. \\ &\sim N(\Gamma_n^{-1} \tilde{\mathbf{y}}_n, \Gamma_n^{-1}) \end{aligned}$$

Approximate likelihood:

$$\begin{aligned} p_a(\mathbf{y}_n; \boldsymbol{\psi}) &= \frac{p(\mathbf{y}_n, \boldsymbol{\alpha}_n)}{p_a(\boldsymbol{\alpha}_n | \mathbf{y}_n)} \propto \frac{\det(G_n)^{1/2}}{\det(\Gamma_n)^{1/2}} \exp\{\mathbf{y}_n^T \mathbf{X}\boldsymbol{\beta} + .5 \tilde{\mathbf{y}}_n^T \Gamma_n^{-1} \tilde{\mathbf{y}}_n\}, \\ \tilde{\mathbf{y}}_n &= \mathbf{y}_n - \exp\{\mathbf{X}\boldsymbol{\beta}\} \exp\{\boldsymbol{\alpha}_n^*\} + \exp\{\boldsymbol{\alpha}_n^*\} \exp\{\mathbf{X}\boldsymbol{\beta}\} \boldsymbol{\alpha}_n^* \end{aligned}$$

(component-wise multiplication for vectors)

Note: We actually expand the joint density for \mathbf{Y}_n and $\boldsymbol{\alpha}_n$ in a neighborhood of $\boldsymbol{\alpha}^*$.

Estimation Methods — Approximation to the likelihood

Implementation:

1. Let $\boldsymbol{\alpha}^* = \boldsymbol{\alpha}^*(\boldsymbol{\psi})$ be the converged value of $\boldsymbol{\alpha}^{(j)}(\boldsymbol{\psi})$, where

$$\boldsymbol{\alpha}^{(j+1)}(\boldsymbol{\psi}) = \Gamma_n^{-1} \tilde{\mathbf{y}}_n(\boldsymbol{\psi})$$

2. Maximize $p_a(\mathbf{y}_n; \boldsymbol{\psi})$ with respect to $\boldsymbol{\psi}$.

Simulation Results

Model: $Y_t | \alpha_t \sim \text{Pois}(\exp(.7 + \alpha_t))$, $\alpha_t = .5 \alpha_{t-1} + \varepsilon_t$, $\{\varepsilon_t\} \sim \text{IID } N(0, .3)$, $n = 200$

Estimation methods:

- Importance sampling (N=1000, ψ_0 updated a maximum of 10 times)


	beta	phi	sigma2
mean	0.6982	0.4718	0.3008
std	0.1059	0.1476	0.0899

- Approximation to likelihood


	beta	phi	sigma2
mean	0.7036	0.4579	0.2962
std	0.0951	0.1365	0.0784

Model: $Y_t | \alpha_t \sim \text{Pois}(\exp(.7 + \alpha_t))$, $\alpha_t = .5 \alpha_{t-1} + \varepsilon_t$, $\{\varepsilon_t\} \sim \text{IID } N(0, .3)$, $n = 200$

Approx likelihood



Importance Sampling



Application to Model Fitting for the Polio Data



Model for $\{\alpha_t\}$:

$$\alpha_t = \phi\alpha_{t-1} + \varepsilon_t, \quad \{\varepsilon_t\} \sim \text{IID } N(0, \sigma^2).$$


- Importance sampling (ψ_0 updated 5 times for each $N=100, 500, 1000,$)
- Simulation based on **1000** replications and the **fitted AL model**.

	Import Sampling			Approx Like			GLM	
	$\hat{\beta}_{IS}$	Simulation		$\hat{\beta}_{AL}$	Simulation		$\hat{\beta}_{GLM}$	SD
		Mean	SD		Mean	SD		
Intercept	0.203	0.223	0.381	0.202	0.210	0.343	.207	0.078
Trend($\times 10^{-3}$)	-2.675	-2.778	3.979	-2.690	-2.720	3.415	-4.18	1.400
$\cos(2\pi t/12)$	0.110	0.103	0.124	0.113	0.111	0.123	-.152	0.097
$\sin(2\pi t/12)$	-0.456	-0.456	0.151	-0.454	-0.454	0.143	-.532	0.109
$\cos(2\pi t/6)$	0.399	0.401	0.123	0.396	0.400	0.114	.169	0.098
$\sin(2\pi t/6)$	0.015	0.024	0.118	0.016	0.012	0.110	-.432	0.101
ϕ	0.865	0.777	0.198	0.845	0.764	0.165		
σ^2	0.088	0.100	0.068	0.104	0.114	0.075		

Application to Model Fitting for the Polio Data (cont)



Polio Data: observed and conditional mean (approx like)



Application to Sydney Asthma Count Data

Data: Y_1, \dots, Y_{1461} daily asthma presentations in a Campbelltown hospital.

Preliminary analysis identified.

- no upward or downward trend
- **annual cycle** modeled by $\cos(2\pi t/365)$, $\sin(2\pi t/365)$
- **seasonal effect** modeled by

$$P_{ij}(t) = \frac{1}{B(2.5,5)} \left(\frac{t - T_{ij}}{100} \right)^{2.5} \left(1 - \frac{t - T_{ij}}{100} \right)^5$$

where $B(2.5,5)$ is the beta function and T_{ij} is the start of the j^{th} school term in year i .

- day of the week effect modeled by separate indicator variables for **Sunday** and **Monday** (increase in admittance on these days compared to Tues-Sat).
- Of the meteorological variables (max/min temp, humidity) and pollution variables (ozone, NO, NO₂), only **humidity** at lags of 12-20 days and **NO₂(max)** appear to have an association.


Results for Asthma Data—(IS & AL)

Term	IS	AL	Mean	SD
Intercept	0.590	0.591	0.593	.0658
Sunday effect	0.138	0.138	0.139	.0531
Monday effect	0.229	0.231	0.230	.0495
$\cos(2\pi t/365)$	-0.218	-0.218	-0.217	.0415
$\sin(2\pi t/365)$	0.200	0.179	0.181	.0437
Term 1, 1990	0.188	0.198	0.194	.0638
Term 2, 1990	0.183	0.130	0.129	.0664
Term 1, 1991	0.080	0.075	0.070	.0733
Term 2, 1991	0.177	0.164	0.157	.0665
Term 1, 1992	0.223	0.221	0.214	.0667
Term 2, 1992	0.243	0.239	0.237	.0620
Term 1, 1993	0.379	0.397	0.394	.0625
Term 2, 1993	0.127	0.111	0.108	.0682
Humidity $H_t/20$	0.009	0.010	0.007	.0032
NO_2 max	-0.125	-0.107	-0.108	.0347
AR(1), ϕ	0.385	0.788	0.468	.3790
σ^2	0.053	0.010	0.018	.0153


Asthma Data: observed and conditional mean

— cond mean
— observed


1990




1991



1992



1993



Summary Remarks

1. Importance sampling offers a nice clean method for estimation in parameter driven models.
2. The innovations algorithm allows for quick implementation of importance sampling. Extends easily to higher-order AR structure.
3. Relative likelihood approach is a one-sample based procedure.
4. Approximation to the likelihood is a non-simulation based procedure which may have great potential especially with large sample sizes and/or large number of explanatory variables. .