

FORMULAS FOR STOPPED DIFFUSION PROCESSES WITH STOPPING TIMES BASED ON DRAWDOWNS AND DRAWUPS

BY LIBOR POSPISIL, JAN VECER, AND OLYMPIA HADJILIADIS,

Columbia University and Brooklyn College (C.U.N.Y.)

This paper studies drawdown and drawup processes in a general diffusion model. The main result is a formula for the joint distribution of the running minimum and the running maximum of the process stopped at the time of the first drop of size a . As a consequence, we obtain the probabilities that a drawdown of size a precedes a drawup of size b and vice versa. The results are applied to several examples of diffusion processes, such as drifted Brownian motion, Ornstein-Uhlenbeck process, and Cox-Ingersoll-Ross process.

1. Introduction. In this article, we study properties of a general diffusion process $\{X_t\}$ stopped at the first time when its drawdown attains a certain value a . Let us denote this time as $T_D(a)$. The drawdown of a process is defined as the current drop of the process from its running maximum. We present two main results here. First, we derive the joint distribution of the running minimum and the running maximum stopped at $T_D(a)$. Second, we calculate the probability that a drawdown of size a precedes a drawup of size b , where the drawup is defined as the increase of $\{X_t\}$ over the running minimum. All formulas are expressed in terms of the drift function, the volatility function, and the initial value of $\{X_t\}$. In addition to the main theorems, this paper contains other results that help us to understand the behavior of diffusion processes better. For example, we relate the probability that the drawup process stopped at $T_D(a)$ is zero to the expected running minimum stopped at $T_D(a)$.

We apply the results to several examples of diffusion processes: drifted Brownian motion, Ornstein-Uhlenbeck process (OU), and Cox-Ingersoll-Ross process (CIR). These examples play important roles in change point detection and in finance. We also discuss how the results presented in this paper are related to the problem of quickest detection and identification of two-sided changes in the drift of general diffusion processes.

Our results extend several theorems stated and proved in Gihman and Skorokhod (1972), Taylor (1975), and Lehoczký (1977). These results include the distribution of a diffusion process stopped at the first time it hits either

AMS 2000 subject classifications: Primary 91B28; secondary 60G44

Keywords and phrases: Diffusion Process, Drawdowns and Drawups, Stopping Time

a lower or an upper barrier, and the distribution of the running maximum of a diffusion process stopped at time $T_D(a)$. The formulas for a drifted Brownian motion presented here coincide with the results in Hadjiliadis and Vecer (2006). The approach used in Hadjiliadis and Vecer (2006) is based on a calculation of the expected first passage times of the drawdown and drawup processes to levels a and b . However, while this approach applies to a drifted Brownian motion, it cannot be extended to a general diffusion process. In this paper, we derive the joint distribution of the running maximum and minimum stopped at $T_D(a)$, which can be obtained for a general diffusion process. Subsequently, we use this result to calculate the probability that a drawdown precedes a drawup.

Properties of drawdown and drawup processes are of interest in change point detection, where the goal is to test whether an abrupt change in a parameter of a dynamical system has occurred. Drawdowns and drawups of the likelihood ratio process serve as test statistics for hypotheses about the change point. Details can be found, for example, in Poor and Hadjiliadis (2008), Hadjiliadis and Moustakides (2006) and Khan (2008).

The concept of a drawdown has been also been studied in applied probability and in finance. The underlying diffusion process usually represents a stock index, an exchange rate, or an interest rate. Some characteristics of its drawdown, such as the expected maximum drawdown, can be used to measure the downside risks of the corresponding market. The distribution of the maximum drawdown of a drifted Brownian motion was determined in Magdon-Ismail et al. (2004). Cherny and Dupire (2007) derived the distribution of a local martingale and its maximum at the first time when the corresponding range process attains value a . Salminen and Vallois (2007) derived the joint distribution of the maximum drawdown and the maximum drawup of a Brownian motion up to an independent exponential time. Vecer (2006) related the expected maximum drawdown of a market to directional trading. Several authors, such as Grossman and Zhou (1993), Cvitanic and Karatzas (1995), and Chekhlov et al. (2005), discussed the problem of portfolio optimization with drawdown constraints. Meilijson (2003) used stopping time $T_D(a)$ to solve an optimal stopping problem based on a drifted Brownian motion and its running maximum. Obloj and Yor (2006) studied properties of martingales with representation $H(M_t, \bar{M}_t)$, where M_t is a continuous local martingale and \bar{M}_t its supremum up to time t . Nikeghbali (2006) associated the Skorokhod's stopping problem with a class of submartingales which includes drawdown processes of continuous local martingales.

This paper is structured in the following way: notation and assumptions are introduced in Section 2. In Section 3, we derive the joint distribution of

the running maximum and the running minimum stopped at the first time that the process drops by a certain amount (Theorem 3.1 and Theorem 3.2), and in Section 4, we calculate the probability that a drawdown of size a will precede a drawup of size b (Theorem 4.1 and Theorem 4.2). Special cases, such as drifted Brownian motion, Ornstein-Uhlenbeck process, Cox-Ingersoll-Ross process, are discussed in Section 5. The relevance of the result in Section 4 to the problem of quickest detection and identification of two-sided alternatives in the drift of general diffusion processes is also presented in Section 5. Finally, Section 6 contains concluding remarks.

2. Drawdown and Drawup Processes. In this section, we define drawdown and drawup processes in a diffusion model and present the main assumptions.

Consider an interval $I = (l, r)$, where $-\infty \leq l < r \leq \infty$. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\{W_t\}$ a Brownian motion, and $\{X_t\}$ a unique strong solution of the following stochastic differential equation:

$$(1) \quad dX_t = \mu(X_t)dt + \sigma(X_t)dW_t, \quad X_0 = x \in I,$$

where $X_t \in I$ for all $t \geq 0$. Moreover, we will assume that functions $\mu(\cdot)$ and $\sigma(\cdot)$ meet the following conditions:

$$(2) \quad \sigma(y) > 0, \quad \text{for } \forall y \in I,$$

$$(3) \quad \int_x^r \frac{\Psi(x, z)}{\int_{z-a}^z \Psi(x, y)dy} dz = \infty, \quad \text{for all } a > 0 \text{ such that } x - a \in I,$$

$$(4) \quad \int_l^x \frac{\Psi(x, z)}{\int_z^{z+b} \Psi(x, y)dy} dz = \infty, \quad \text{for all } b > 0 \text{ such that } x + b \in I,$$

where $\Psi(u, z) = e^{-2 \int_u^z \gamma(y)dy}$ and $\gamma(y) = \frac{\mu(y)}{\sigma^2(y)}$. Drifted Brownian motion, Ornstein-Uhlenbeck process, and Cox-Ingersoll-Ross process are examples of diffusion processes satisfying these assumptions. Functions $\mu(\cdot)$ and $\sigma(\cdot)$ will be referred to as the drift and the volatility functions. Note that a process given by (1) has the strong Markov property. If we need to emphasize that x is the starting value of $\{X_t\}$, we will write $\mathbb{P}_x[\cdot]$.

Let us define the running maximum, $\{M_t\}$, and the running minimum, $\{m_t\}$, of process $\{X_t\}$ as:

$$M_t = \sup_{s \in [0, t]} X_s, \quad m_t = \inf_{s \in [0, t]} X_s.$$

The drawdown and the drawup of $\{X_t\}$ are defined as:

$$DD_t = M_t - X_t, \quad DU_t = X_t - m_t.$$

We denote by $T_D(a)$ and $T_U(b)$ the first passage times of the processes $\{DD_t\}$ and $\{DU_t\}$ to the levels a and b respectively, where $a > 0$, $b > 0$, $x - a \in I$, and $x + b \in I$. We set $T_D(a) = \infty$ or $T_U(b) = \infty$ if process DD_t does not reach a or process DU_t does not reach b :

$$T_D(a) = \inf\{t \geq 0; DD_t = a\}, \quad T_U(b) = \inf\{t \geq 0; DU_t = b\}.$$

Conditions (3) and (4) ensure that

$$\begin{aligned} \mathbb{P}_x[T_D(a) < \infty] &= \lim_{v \rightarrow r-} \mathbb{P}_x[M_{T_D(a)} \leq v] = 1, \\ \mathbb{P}_x[T_U(b) < \infty] &= \lim_{u \rightarrow l+} \mathbb{P}_x[m_{T_U(b)} > u] = 1, \end{aligned}$$

Thus, we assume that $T_D(a) < \infty$ and $T_U(b) < \infty$ almost surely for any $a > 0$ and $b > 0$, such that $x - a \in I$ and $x + b \in I$.

In the following sections, we derive the joint distribution of $(m_{T_D(a)}, M_{T_D(a)})$ (Section 3) and a formula for the probability $\mathbb{P}_x[T_D(a) < T_U(b)]$ (Section 4).

3. Joint Distribution of the Running Minimum and the Running Maximum Stopped at $T_D(a)$. The distribution of random variable $M_{T_D(a)}$ was derived in Lehoczky (1977). In our paper, we focus on the joint distribution of $(m_{T_D(a)}, M_{T_D(a)})$.

Note that the running minimum stopped at time $T_D(a)$ is bounded by $x - a$ and x : $x - a \leq m_{T_D(a)} \leq x$. The joint distribution of the running minimum and maximum stopped at time $T_D(a)$ will be denoted as \bar{H} :

$$\bar{H}_x(u, v) = \mathbb{P}_x[m_{T_D(a)} > u, M_{T_D(a)} > v],$$

where $u \in [x - a, x]$ and $v \in [x, \infty)$. In the following theorem, we will express \bar{H} in terms of function $\Psi(u, z) = e^{-2 \int_u^z \gamma(y) dy}$, where $\gamma(y) = \frac{\mu(y)}{\sigma^2(y)}$.

Theorem 3.1 *Let $a > 0$ such that $x - a \in I$. The random variables $m_{T_D(a)}$ and $M_{T_D(a)}$ have the following joint distribution:*

$$(5) \quad \bar{H}_x(u, v) = \frac{\int_u^x \Psi(u, z) dz}{\int_u^{u+a} \Psi(u, z) dz} e^{-\int_{u+a}^v \frac{\Psi(u+a, z)}{\int_{z-a}^z \Psi(u+a, y) dy} dz},$$

where $u \in [x - a, x]$, $v \in [u + a, \infty)$, $\Psi(u, z) = e^{-2 \int_u^z \gamma(y) dy}$ and $\gamma(y) = \frac{\mu(y)}{\sigma^2(y)}$. If $u \in [x - a, x]$ and $v \in [x, u + a)$, then:

$$(6) \quad \bar{H}_x(u, v) = \frac{\int_u^x \Psi(u, z) dz}{\int_u^{u+a} \Psi(u, z) dz}.$$

PROOF: The process $\{X_t\}$ is given by (1) and $X_0 = x$. First, let us assume that $u \in [x - a, x]$ and $v \in [u + a, \infty)$. The event $\{m_{T_D(a)} > u, M_{T_D(a)} > v\}$ occurs if and only if the process $\{X_t\}$ attains $u + a$ without dropping below u and then exceeds v before the drawdown reaches a . Due to the Markov property of the process $\{X_t\}$, we can write the probabilities of these events as follows:

$$\begin{aligned}
 \bar{H}_x(u, v) &= \mathbb{P}_x[m_{T_D(a)} > u, M_{T_D(a)} > v] \\
 &= \mathbb{P}_x[X_{\tau(u, u+a)} = u + a] \mathbb{P}_{u+a}[M_{T_D(a)} > v] \\
 (7) \quad &= \frac{\int_u^x \Psi(u, z) dz}{\int_u^{u+a} \Psi(u, z) dz} e^{-\int_{u+a}^v \frac{\Psi(u+a, z)}{\int_{z-a}^z \Psi(u+a, y) dy} dz},
 \end{aligned}$$

where $\tau(u, u + a) = \inf\{t \geq 0; X_t = u \text{ or } X_t = u + a\}$. The formula for the first probability in (7) follows from Gihman and Skorokhod (1972), page 110. The second probability in (7), representing the survival function of $M_{T_D(a)}$, was derived in Lehoczky (1977), page 602. Finally, if $v < u + a$, we have $\{m_{T_D(a)} > u, M_{T_D(a)} > v\} = \{m_{T_D(a)} > u\} = \{X_{\tau(u, u+a)} = u + a\}$ because $M_{T_D(a)} \geq m_{T_D(a)} + a$. Thus,

$$\bar{H}_x(u, v) = \frac{\int_u^x \Psi(u, z) dz}{\int_u^{u+a} \Psi(u, z) dz}.$$

◇

The distribution function, the survival function, and the density function of $m_{T_D(a)}$ will be denoted as F , \bar{F} , and f , respectively:

$$\bar{F}_x(u) = 1 - F_x(u) = \mathbb{P}_x[m_{T_D(a)} > u], \quad f_x(u) = \frac{dF_x(u)}{du},$$

where $u \in [x - a, x]$. We can derive the marginal distribution of $m_{T_D(a)}$ from the results in Theorem 3.1.

Corollary 3.2 *Let $a > 0$ such that $x - a \in I$. The distribution function, the density function, and the expected value of random variable $m_{T_D(a)}$ are:*

$$(8) \quad \bar{F}_x(u) = \frac{\int_u^x \Psi(u, z) dz}{\int_u^{u+a} \Psi(u, z) dz},$$

$$(9) \quad f_x(u) = \frac{\Psi(u, u+a) \int_u^x \Psi(u, z) dz + \int_x^{u+a} \Psi(u, z) dz}{\left(\int_u^{u+a} \Psi(u, z) dz\right)^2},$$

$$(10) \quad \mathbb{E}_x \left[m_{T_D(a)} \right] = x - \int_{x-a}^x \frac{\int_x^{u+a} \Psi(u, z) dz}{\int_u^{u+a} \Psi(u, z) dz} du,$$

where $u \in [x-a, x]$, $\Psi(u, z) = e^{-2 \int_u^z \gamma(y) dy}$, and $\gamma(y) = \frac{\mu(y)}{\sigma^2(y)}$.

Let us denote the distribution function and the survival function of the running maximum stopped at $T_D(a)$ as G and \bar{G} :

$$\bar{G}_x(v) = 1 - G_x(v) = \mathbb{P}_x[M_{T_D(a)} > v], \quad x \leq v < r.$$

Note that the joint distribution of $(m_{T_D(a)}, M_{T_D(a)})$ can be represented by the marginal distributions \bar{F} and \bar{G} :

$$(11) \quad \bar{H}_x(u, v) = \bar{F}_x(u) \bar{G}_{u+a}(v),$$

where $u \in [x-a, x]$ and $v \in [u+a, \infty)$. Let us calculate the derivative of $\bar{H}_x(u, v)$ with respect to u , which will be used in the proof of the main theorem.

Lemma 3.3 *Let $a > 0$ such that $x-a \in I$. Let $u \in [x-a, x]$ and $v \geq u+a$. Then*

$$(12) \quad -\frac{\partial \bar{H}_x}{\partial u}(u, v) = e^{-\int_{u+a}^v \frac{\Psi(u+a, z)}{\int_{z-a}^z \Psi(u+a, y) dy} dz} \frac{\int_x^{u+a} \Psi(u, z) dz}{\left(\int_u^{u+a} \Psi(u, z) dz \right)^2},$$

where $\Psi(u, z) = e^{-2 \int_u^z \gamma(y) dy}$ and $\gamma(y) = \frac{\mu(y)}{\sigma^2(y)}$.

PROOF: According to (11),

$$\frac{\partial \bar{H}_x}{\partial u}(u, v) = \frac{\partial \bar{F}_x(u)}{\partial u} \bar{G}_{u+a}(v) + \bar{F}_x(u) \frac{\partial \bar{G}_{u+a}(v)}{\partial u}.$$

Note that the function Ψ has the following property: $\Psi(a, b)\Psi(b, c) = \Psi(a, c)$.

Therefore, $\frac{\int \Psi(u, z) dz}{\int \Psi(u, z) dz} = \frac{\int \Psi(C, z) dz}{\int \Psi(C, z) dz}$ and $\frac{\int \Psi(u, y)}{\int \Psi(u, z) dz} = \frac{\int \Psi(C, y)}{\int \Psi(C, z) dz}$ for any constant C . Thus, the first variable of Ψ is redundant in such fractions and can be omitted during the calculation of their derivative with respect to u . Using formula (9), we have

$$\frac{\partial \bar{F}_x(u)}{\partial u} = -f_x(u) = \frac{-\int_x^{u+a} \Psi(u, z) dz - \Psi(u, u+a) \int_u^x \Psi(u, z) dz}{\left(\int_u^{u+a} \Psi(u, z) dz \right)^2}.$$

The derivative of $\bar{G}_{u+a}(v)$ with respect to u is given by:

$$\begin{aligned} \frac{\partial \bar{G}_{u+a}(v)}{\partial u} &= \frac{1}{\int_u^{u+a} \Psi(u+a, z) dz} e^{-\int_x^{u+a} \frac{\Psi(u+a, z)}{\int_{z-a}^z \Psi(u+a, y) dy} dz} \\ &= \frac{\Psi(u, u+a)}{\int_u^{u+a} \Psi(u, z) dz} \bar{G}_{u+a}(v). \end{aligned}$$

Combining these results yields formula (12). \diamond

Formula (9) allows us to decompose the density of $m_{T_D(a)}$ into two parts:

$$(13) \quad f_x(u)du = \mathbb{P}_x[DU_{T_D(a)} > 0, m_{T_D(a)} \in (u, u + du)] \\ + \mathbb{P}_x[DU_{T_D(a)} = 0, m_{T_D(a)} \in (u, u + du)].$$

The set $\{DU_{T_D(a)} = 0\}$ corresponds to the event that the process attained its running minimum at time $T_D(a) : X_{T_D(a)} = m_{T_D(a)}$. In the following lemma, we calculate the probabilities introduced in (13).

Lemma 3.4 *Let $a > 0$ such that $x - a \in I$. Then*

$$(14) \quad \mathbb{P}_x[DU_{T_D(a)} > 0, m_{T_D(a)} \in (u, u + du)] = \frac{\int_x^{u+a} \Psi(u, z) dz}{\left(\int_u^{u+a} \Psi(u, z) dz\right)^2} du,$$

$$(15) \quad \mathbb{P}_x[DU_{T_D(a)} = 0, m_{T_D(a)} \in (u, u + du)] = \frac{\Psi(u, u + a) \int_u^x \Psi(u, z) dz}{\left(\int_u^{u+a} \Psi(u, z) dz\right)^2} du,$$

$$(16) \quad \mathbb{P}_x[DU_{T_D(a)} = 0] = \int_{x-a}^x \frac{\Psi(u, u + a) \int_u^x \Psi(u, z) dz}{\left(\int_u^{u+a} \Psi(u, z) dz\right)^2} du,$$

where $\Psi(u, z) = e^{-2 \int_u^z \gamma(y) dy}$ and $\gamma(y) = \frac{\mu(y)}{\sigma^2(y)}$.

PROOF: Let us use the relationship $M_{T_D(a)} = m_{T_D(a)} + DU_{T_D(a)} + a$ to rewrite probability (14) in terms of the function $\overline{H}_x(u, v)$:

$$\mathbb{P}_x[DU_{T_D(a)} > 0, m_{T_D(a)} \in (u, u + du)] = \\ = \mathbb{P}_x[M_{T_D(a)} > u + a, m_{T_D(a)} \in (u, u + du)] = -\frac{\partial \overline{H}_x}{\partial u}(u, u + a) du$$

for $u \in [x - a, x]$. Thus, result (14) follows from Lemma 3.3. Formula (15) can be obtained using the decomposition of f in (13) and equations (9) and (14). The result (15) also implies (16):

$$\mathbb{P}_x[DU_{T_D(a)} = 0] = \int_{x-a}^x \mathbb{P}_x[DU_{T_D(a)} = 0, m_{T_D(a)} \in (u, u + du)].$$

\diamond

Remark 3.5 *If $a > 0$ such that $x - a \in I$,*

$$(17) \quad \mathbb{P}_x[DU_{T_D(a)} = 0] = -\frac{\partial}{\partial a} \mathbb{E}_x[m_{T_D(a)}].$$

PROOF: Formula (17) can be verified by calculating the derivative of $\mathbb{E}_x[m_{T_D(a)}]$, which is given by (10), and comparing the result with (16). \diamond

Let us heuristically explain Remark 3.5 using the following expression:

$$m_{T_D(a)} = (M_{T_D(a)} - a) \mathbb{I}_{\{DU_{T_D(a)}=0\}} + m_{T_D(a)} \mathbb{I}_{\{DU_{T_D(a)}>0\}}.$$

Shifting a by a small number h has an impact on $m_{T_D(a)}$ only if the process $\{X_t\}$ attained its running minimum m at time $T_D(a)$:

$$DU_{T_D(a)} = X_{T_D(a)} - m_{T_D(a)} = 0.$$

In this case, $m_{T_D(a)} = M_{T_D(a)} - a$ and the change in $m_{T_D(a)}$ is $-h$ because the running maximum $M_{T_D(a)}$ remains the same. On the other hand, if $\{X_t\}$ is greater than $m_{T_D(a)}$, that is, $DU_{T_D(a)} > 0$, then small changes in a do not affect $m_{T_D(a)}$. As a result,

$$m_{T_D(a+h)} - m_{T_D(a)} \approx -h \mathbb{I}_{\{DU_{T_D(a)}=0\}}$$

for h small. Applying the expected value to this relationship and letting $h \rightarrow 0$ leads to the equation (17).

The knowledge of the joint distribution $\bar{H}_x(u, v)$ allows us to determine the distribution and the expected value of the range process, $R_t = M_t - m_t$, stopped at time $T_D(a)$.

Corollary 3.6 *Let $a > 0$ such that $x - a \in I$. The distribution of the range process $R_t = M_t - m_t$, stopped at time $T_D(a)$ is:*

$$(18) \quad \mathbb{P}_x[R_{T_D(a)} > r] = \int_{x-a}^x \bar{G}_{u+a}(u+r) \frac{\int_x^{u+a} \Psi(u,z) dz}{\left(\int_u^{u+a} \Psi(u,z) dz\right)^2} du,$$

$$(19) \quad \mathbb{P}_x[R_{T_D(a)} = a] = \int_{x-a}^x \frac{\Psi(u, u+a) \int_u^x \Psi(u,z) dz}{\left(\int_u^{u+a} \Psi(u,z) dz\right)^2} du,$$

where $r > a$. The expected value of $R_{T_D(a)}$:

$$(20) \quad \mathbb{E}_x[R_{T_D(a)}] = \int_x^\infty \bar{G}_x(v) dv + \int_{x-a}^x \frac{\int_x^{u+a} \Psi(u,z) dz}{\int_u^{u+a} \Psi(u,z) dz} du,$$

where

$$\begin{aligned}\bar{G}_{u+a}(v) &= e^{-\int_{u+a}^v \frac{\Psi(u+a,z)}{\int_{z-a}^z \Psi(u+a,y)dy} dz}, \\ \Psi(u,z) &= e^{-2\int_u^z \gamma(y)dy}, \quad \text{and} \quad \gamma(y) = \frac{\mu(y)}{\sigma^2(y)}.\end{aligned}$$

4. Probability of a Drawdown Preceding a Drawup. In this section, we derive formulas for the probabilities that a drawdown of size a precedes a drawup of size b and vice versa. The calculation is based on the knowledge of the joint distribution of $(m_{T_D(a)}, M_{T_D(a)})$, which appears in Theorem 3.1.

Theorem 4.1 *Assume that $\{X_t\}$ is a unique strong solution of equation (1) and that conditions (2), (3), and (4) are satisfied. Let $b \geq a > 0$ such that $x - a \in I$ and $x + b \in I$. Then:*

$$(21) \quad \mathbb{P}_x[T_D(a) < T_U(b)] = 1 - \int_{x-a}^x \bar{G}_{u+a}(u+b) \frac{\int_x^{u+a} \Psi(u,z)dz}{\left(\int_u^{u+a} \Psi(u,z)dz\right)^2} du,$$

$$(22) \quad \mathbb{P}_x[T_D(a) > T_U(b)] = \int_{x-a}^x \bar{G}_{u+a}(u+b) \frac{\int_x^{u+a} \Psi(u,z)dz}{\left(\int_u^{u+a} \Psi(u,z)dz\right)^2} du,$$

where

$$\begin{aligned}\bar{G}_{u+a}(v) &= e^{-\int_{u+a}^v \frac{\Psi(u+a,z)}{\int_{z-a}^z \Psi(u+a,y)dy} dz}, \\ \Psi(u,z) &= e^{-2\int_u^z \gamma(y)dy}, \quad \text{and} \quad \gamma(y) = \frac{\mu(y)}{\sigma^2(y)}.\end{aligned}$$

PROOF: Let $b \geq a > 0$. First, we will show that

$$(23) \quad \{T_D(a) < T_U(b)\} = \{0 < DU_{T_D(a)} < b - a\} \cup \{DU_{T_D(a)} = 0\}.$$

The range of the process $\{X_t\}$ at t is defined as $R_t = M_t - m_t$, which implies $R_t = DU_t + DD_t$. One can also prove that

$$(24) \quad R_t = \max\left\{\sup_{[0,t]} DU_u, \sup_{[0,t]} DD_u\right\}.$$

The process on the right hand side of (24) is non-decreasing and equals zero at time $t = 0$. Moreover, it increases only if $\sup_{[0,t]} DU_u$ or $\sup_{[0,t]} DD_u$ changes, which occurs when either $X_t = M_t$ or $X_t = m_t$. In this case, the right hand side is $M_t - m_t$, which justifies (24).

According to the formula (24), $DU_{T_D(a)} + a = \max\{\sup_{[0,T_D(a)]} DU_u, a\}$. Thus, $DU_{T_D(a)} = \max\{\sup_{[0,T_D(a)]} DU_u - a, 0\}$ and

$$\begin{aligned} \{T_D(a) < T_U(b)\} &= \left\{ \sup_{[0,T_D(a)]} DU_u < b \right\} \\ &= \{0 < DU_{T_D(a)} < b - a\} \cup \{DU_{T_D(a)} = 0\}, \end{aligned}$$

which proves (23). Furthermore, using the relationship $M_{T_D(a)} = m_{T_D(a)} + DU_{T_D(a)} + a$, we have:

$$\begin{aligned} \mathbb{P}_x[DU_{T_D(a)} > b - a, m_{T_D(a)} \in (u, u + du)] &= \\ &= \mathbb{P}_x[M_{T_D(a)} > u + b, m_{T_D(a)} \in (u, u + du)] = -\frac{\partial \bar{H}_x}{\partial u}(u, u + b)du. \end{aligned}$$

Now let us calculate the probability of the event $\{T_D(a) < T_U(b)\}$:

$$\begin{aligned} \mathbb{P}_x[T_D(a) < T_U(b)] &= \mathbb{P}_x[DU_{T_D(a)} = 0] + \mathbb{P}_x[0 < DU_{T_D(a)} < b - a] \\ &= 1 - \mathbb{P}_x[DU_{T_D(a)} > b - a] \\ &= 1 - \int_{x-a}^x \mathbb{P}_x[DU_{T_D(a)} > b - a, m_{T_D(a)} \in (u, u + du)] \\ &= 1 - \int_{x-a}^x \mathbb{P}_x[M_{T_D(a)} > u + b, m_{T_D(a)} \in (u, u + du)] \\ (25) \quad &= 1 - \int_{x-a}^x \left\{ -\frac{\partial \bar{H}_x}{\partial u}(u, u + b) \right\} du. \end{aligned}$$

The derivative of \bar{H} is calculated in Lemma 3.3. If we replace $\left\{ -\frac{\partial \bar{H}_x}{\partial u}(u, u + b) \right\}$ in (25) with that result, we obtain formula (21). Probability (22) is the complement of (21). \diamond

If $b < a$, the formula for $\mathbb{P}_x[T_D(a) < T_U(b)]$ is a modification of (21).

Theorem 4.2 *Assume that $\{X_t\}$ is a unique strong solution of equation (1) and that conditions (2), (3), and (4) are satisfied. Let $0 < b < a$ such that $x - a \in I$ and $x + b \in I$. Then:*

$$(26) \quad \mathbb{P}_x[T_D(a) < T_U(b)] = \int_x^{x+b} G_{v-b}^*(v-a) \frac{\int_{v-b}^x \Psi(v,z) dz}{\left(\int_{v-b}^v \Psi(v,z) dz \right)^2} dv,$$

$$(27) \quad \mathbb{P}_x[T_D(a) > T_U(b)] = 1 - \int_x^{x+b} G_{v-b}^*(v-a) \frac{\int_{v-b}^x \Psi(v,z) dz}{\left(\int_{v-b}^v \Psi(v,z) dz\right)^2} dv,$$

where

$$G_{v-b}^*(u) = e^{-\int_u^{v-b} \frac{\Psi(v-b,z)}{\int_z^{z+b} \Psi(v-b,y) dy} dz},$$

$$\Psi(u,z) = e^{-2\int_u^z \gamma(y) dy}, \quad \text{and} \quad \gamma(y) = \frac{\mu(y)}{\sigma^2(y)}.$$

PROOF: The proof is analogous to the proof of Theorem 4.1. \diamond

The procedure we used in the proof of Theorem 4.1 allows us to interpret equations (21) and (22) as follows:

$$\begin{aligned} \mathbb{P}_x[T_D(a) < T_U(b)] &= 1 - \int_{x-a}^x \mathbb{P}_x[M_{T_D(a)} > u+b, m_{T_D(a)} \in (u, u+du)] \\ &= \int_{x-a}^x \mathbb{P}_x[M_{T_D(a)} \leq u+b, m_{T_D(a)} \in (u, u+du)], \end{aligned}$$

$$\mathbb{P}_x[T_D(a) > T_U(b)] = \int_{x-a}^x \mathbb{P}_x[M_{T_D(a)} > u+b, m_{T_D(a)} \in (u, u+du)].$$

Let us discuss this interpretation. If $m_{T_D(a)}$ lies in a neighborhood of u , the event of $\{T_D(a) > T_U(b)\}$ coincides with $\{M_{T_D(a)} > u+b\}$. Probability $\mathbb{P}_x[T_D(a) > T_U(b)]$ is then the integral of $\mathbb{P}_x[M_{T_D(a)} > u+b, m_{T_D(a)} \in (u, u+du)]$ over all possible values of $m_{T_D(a)}$.

When $a = b$ in (21) and (22), we have $\{T_D(a) < T_U(a)\} = \{DU_{T_D(a)} = 0\}$:

$$\mathbb{P}_x[T_D(a) < T_U(a)] = \int_{x-a}^x \frac{\Psi(u, u+a) \int_u^x \Psi(u,z) dz}{\left(\int_u^{u+a} \Psi(u,z) dz\right)^2} du,$$

$$\mathbb{P}_x[T_D(a) > T_U(a)] = \int_{x-a}^x \frac{\int_x^{u+a} \Psi(u,z) dz}{\left(\int_u^{u+a} \Psi(u,z) dz\right)^2} du.$$

5. Application of the Results. In this section, we apply the results from Theorem 3.2 and Theorem 4.1 to the following examples of diffusion processes: Brownian motion, Ornstein-Uhlenbeck process, Cox-Ingersoll-Ross process. We also present an application of the result in Theorem 4.1 to the problem of quickest detection and identification of two-sided changes in the drift of general diffusion processes.

Process	I	$\mu(y)$	$\sigma(y)$	$\Psi(u, z)$
Brownian motion	\mathbb{R}	μ	σ	$e^{-\frac{2\mu}{\sigma^2}(z-u)}$
Ornstein-Uhlenbeck process	\mathbb{R}	$\kappa(\theta - y)$	σ	$e^{\frac{\kappa}{\sigma^2}[(z-\theta)^2 - (u-\theta)^2]}$
Cox-Ingersoll-Ross process	$(0, \infty)$	$\kappa(\theta - y)$	$\sigma\sqrt{y}$	$\left(\frac{z}{u}\right)^{-\frac{2\kappa\theta}{\sigma^2}} e^{\frac{2\kappa}{\sigma^2}(z-u)}$

TABLE 1

Function $\Psi(u, z)$ for examples of diffusion processes.

5.1. Examples of Diffusion Processes. The formulas for $\bar{G}_x(v)$, $\bar{F}_x(v)$, $\bar{H}_x(u, v)$, and $\mathbb{P}_x[T_D(a) < T_U(b)]$ depend on function $\Psi(u, z)$. Table 1 shows specific forms of this function for several examples of diffusion processes $dX_t = \mu(X_t)dt + \sigma(X_t)dW_t$, where $X_t \in I$ and $X_0 = x$.

We assume that conditions (2), (3), and (4) are satisfied for all these processes. In the cases of a drifted Brownian motion and an Ornstein-Uhlenbeck process, the conditions hold true for any combination of the parameters. In the case of a Cox-Ingersoll-Ross process, we need to make an additional assumption: $k\theta > \sigma^2/2$.

One can derive an analytical expression of the function $\bar{H}_x(u, v)$ and the probability $\mathbb{P}_x[T_D(a) < T_U(b)]$ for Brownian motion:

$$\bar{H}_x(u, v) = \frac{e^{\frac{2\mu}{\sigma^2}a} - e^{\frac{2\mu}{\sigma^2}(u-(x-a))}}{e^{\frac{2\mu}{\sigma^2}a} - 1} \exp\left\{- (v-x) \frac{\frac{2\mu}{\sigma^2}}{e^{\frac{2\mu}{\sigma^2}a} - 1}\right\},$$

where $u \in [x-a, x]$ and $v \in [u+a, \infty)$, implying

$$\bar{G}_x(v) = \mathbb{P}_x[M_{T_D(a)} > v] = \exp\left\{- (v-x) \frac{\frac{2\mu}{\sigma^2}}{e^{\frac{2\mu}{\sigma^2}a} - 1}\right\}, \quad v \in [x, \infty),$$

$$\bar{F}_x(u) = \mathbb{P}_x[m_{T_D(a)} > u] = \frac{e^{\frac{2\mu}{\sigma^2}a} - e^{\frac{2\mu}{\sigma^2}(u-(x-a))}}{e^{\frac{2\mu}{\sigma^2}a} - 1}, \quad u \in [x-a, x],$$

$$\mathbb{E}_x[m_{T_D(a)}] = x - \frac{\sigma^2}{2\mu} + \frac{a}{e^{\frac{2\mu}{\sigma^2}a} - 1},$$

$$\mathbb{P}_x[T_D(a) < T_U(b)] = 1 - \exp\left\{- (b-a) \frac{\frac{2\mu}{\sigma^2}}{e^{\frac{2\mu}{\sigma^2}a} - 1}\right\} \frac{e^{\frac{2\mu}{\sigma^2}a} - \frac{2\mu}{\sigma^2}a - 1}{e^{\frac{2\mu}{\sigma^2}a} + e^{-\frac{2\mu}{\sigma^2}a} - 2},$$




FIG 1. Densities of $M_{T_D(a)}$ and $m_{T_D(a)}$, where $a = 1$, for a drifted Brownian motion: $X_t = \mu t + \sigma W_t$. The densities depend on the parameters through ratio μ/σ^2 . $M_{T_D(a)}$ has an exponential distribution on $[0, \infty)$, while $m_{T_D(a)}$ has a uniform distribution on $[-1, 0]$ if $\mu = 0$ and a truncated exponential distribution on $[-1, 0]$ otherwise.

where $b \geq a > 0$. If $a = b$,

$$\mathbb{P}_x[T_D(a) < T_U(a)] = \frac{e^{-\frac{2\mu}{\sigma^2}a} + \frac{2\mu}{\sigma^2}a - 1}{e^{\frac{2\mu}{\sigma^2}a} + e^{-\frac{2\mu}{\sigma^2}a} - 2}.$$

Random variable $M_{T_D(a)}$ has an exponential distribution on $[x, \infty)$ and $m_{T_D(a)}$ has a truncated exponential distribution on $[x - a, x]$. Note that the formula for $\mathbb{P}_x[T_D(a) < T_U(b)]$ is identical with the results presented in Hadjiliadis and Vecer (2006).

When the drift μ equals to zero, the formulas further reduce to:

$$\overline{H}_x(u, v) = \mathbb{P}_x[m_{T_D(a)} > u, M_{T_D(a)} > v] = \frac{x - u}{a} e^{-\frac{v-(u+a)}{a}},$$

where $u \in [x - a, x]$ and $v \in [u + a, \infty)$, implying

$$\overline{G}_x(v) = \mathbb{P}_x[M_{T_D(a)} > v] = e^{-\frac{v-x}{a}}, \quad v \in [x, \infty),$$

$$\overline{F}_x(u) = \mathbb{P}_x[m_{T_D(a)} > u] = \frac{x - u}{a} \quad \text{and} \quad \mathbb{E}_x[m_{T_D(a)}] = x - \frac{a}{2},$$




FIG 2. Probability $\mathbb{P}_x[T_D(a) < T_U(b)]$ as a function of μ/σ^2 for different values of a and b . We assume that $\{X_t\}$ is a drifted Brownian motion, $X_t = \mu t + \sigma W_t$. Note that $\mathbb{P}_x[T_D(1) < T_U(1)] = 0.5$ for $\mu = 0$.

$$\mathbb{P}_x[T_D(a) < T_U(b)] = 1 - \frac{1}{2}e^{-\frac{b-a}{a}},$$

where $b \geq a > 0$. If $a = b$,

$$\mathbb{P}_x[T_D(a) < T_U(a)] = \mathbb{P}_x[T_D(a) > T_U(a)] = \frac{1}{2}.$$

Hence, $M_{T_D(a)}$ has an exponential distribution on $[x, \infty)$ with parameter $\frac{1}{a}$ and $m_{T_D(a)}$ has a uniform distribution on $[x - a, x]$.

Calculation of $\mathbb{P}_x[T_D(a) < T_U(b)]$ for an Ornstein-Uhlenbeck process and a Cox-Ingersoll-Ross process involves numerical integration.

In Figures 1, 3, and 5, we have plotted densities of $M_{T_D(a)}$ and $m_{T_D(a)}$ for various diffusion processes. Figures 2, 4, and 6 capture dependence of the probability $\mathbb{P}_x[T_D(a) < T_U(b)]$ on the parameters of the processes.

Let us discuss the interpretation of Figure 4, which shows the probability $\mathbb{P}_x[T_D(1) < T_U(1)]$ as a function of κ/σ^2 . When $\kappa = 0$, the drift term of $\{X_t\}$ vanishes and the probability is $\frac{1}{2}$. Moreover, if the process starts at its long-term mean, $x = \theta$, it is symmetric and $\mathbb{P}_x[T_D(1) < T_U(1)] = \frac{1}{2}$ for any value of κ/σ^2 . Now let us assume that $x = \theta + 1$. As κ/σ^2 increases, the drift term will prevail over the volatility term and the process will be pushed down from x to θ . As a result, a drawdown of size 1 will tend to occur before a drawup of size 1, which explains the convergence of $\mathbb{P}_x[T_D(a) < T_U(a)]$ to 1 as $\kappa/\sigma^2 \rightarrow \infty$. Similar reasoning can be used to justify the convergence of $\mathbb{P}_x[T_D(1) < T_U(1)]$ to 0 if $x = \theta - 1$.




FIG 3. Densities of $M_{T_D^{(a)}}$ and $m_{T_D^{(a)}}$, where $a = 1$, assuming that $\{X_t\}$ is an Ornstein-Uhlenbeck process: $dX_t = \kappa(\theta - X_t)dt + \sigma dW_t$, $X_0 = x$. The densities depend on parameters κ and σ through ratio κ/σ^2 . Note that if $x = \theta$, process $\{X_t\}$ is symmetric and consequently, $m_{T_D^{(a)}}$ has a symmetric distribution.

5.2. The problem of quickest detection and identification. In this example, we present the problem of quickest detection and identification of two-sided changes in the drift of a general diffusion process. More specifically, we give precise calculations of the probability of misidentification of two-sided alternatives. In particular, let $\{X_t\}$ be a diffusion process with the initial value $X_0 = x$ and the following dynamics up to a deterministic time τ :

$$(28) \quad dX_t = \sigma(X_t)dW_t, \quad t \leq \tau.$$

For $t > \tau$, the process evolves according to one of the following stochastic differential equations:

$$(29) \quad dX_t = \mu(X_t)dt + \sigma(X_t)dW_t \quad t > \tau,$$

$$(30) \quad dX_t = -\mu(X_t)dt + \sigma(X_t)dW_t \quad t > \tau.$$

with initial condition $y = X_\tau$. We assume that the functions $\mu(\cdot)$ and $\sigma(\cdot)$ are known and the stochastic differential equations (28), (29), and (30) satisfy conditions (2), (3), and (4) stated in Section 1.




FIG 4. Probability $\mathbb{P}_x[T_D(a) < T_U(b)]$, where $a = b = 1$, as a function of κ/σ^2 . Process $\{X_t\}$ is an Ornstein-Uhlenbeck process: $dX_t = \kappa(\theta - X_t)dt + \sigma dW_t$, $X_0 = x$. If $x = \theta$, then the process is symmetric and the probability is 0.5 for any value of κ/σ^2 .

The time of the regime change, τ , is deterministic but unknown. We observe the process $\{X_t\}$ sequentially and our goal is to identify which regime is in effect after τ .

In this context suppose that the first passage time of the drawup process to a threshold a , $T_U(a)$, can be used as a means of detecting the change of dynamics of $\{X_t\}$ from (28) to (29). Similarly, suppose that the first passage time of the drawdown process to a threshold b , $T_D(b)$ may be used as a means of detecting the change of dynamics of $\{X_t\}$ from (28) to (30) (see Khan (2008), Poor and Hadjiliadis (2008)). The simplest example is when $\mu(X_t) = \mu$.

The probability measures $\mathbb{P}_x^{\tau,(1)}$ and $\mathbb{P}_x^{\tau,(2)}$ are the measures generated on the space of continuous functions $C[0, \infty)$ by the process $\{X_t\}$, if the regime changes at time τ from (28) to (29) and from (28) to (30), respectively. The stopping rule proposed and used widely in the literature for detecting such a change is known as the two-sided CUSUM (Cumulative Sum) test, $T(a) = \min\{T_D(a), T_U(a)\}$. This rule was proposed in 1959 by Barnard. Its properties have been widely studied by many authors (Kemp (1961), van Dobben de Bruyn (1968) Bisell (1969), Woodall (1984), Khan (2008)) and a version of this rule was also proven asymptotically optimal in Hadjiliadis and Moustakides (2006). It is thus the rule that has been established in the literature for detecting two-sided changes in the set-up described above.

Theorem 4.1 can be used to compute the probability of a false identifica-




FIG 5. Densities of $M_{T_D(a)}$ and $m_{T_D(a)}$, where $a = 1$. $\{X_t\}$ is a Cox-Ingersoll-Ross process: $dX_t = \kappa(\theta - X_t)dt + \sigma\sqrt{X_t}dW_t$, $X_0 = x$. We use the same values of parameters as in Figure 3.

tion of the change. More specifically,

$$\begin{aligned}
 \mathbb{P}_x^{0,(1)}[T(a) = T_D(a)] &= \mathbb{P}_x^{0,(1)}[T_D(a) \leq T_U(a)] \\
 (31) \qquad \qquad \qquad &= \int_{x-a}^x \frac{\Psi(u, u+a) \int_u^x \Psi(u, z) dz}{\left(\int_u^{u+a} \Psi(u, z) dz\right)^2} du,
 \end{aligned}$$

with $\Psi(u, z) = e^{-2 \int_u^z \gamma(y) dy}$ and $\gamma(y) = \frac{\mu(y)}{\sigma^2(y)}$, expresses the probability that an alarm indicating that the regime switched to (30) will occur before an alarm indicating that the regime switched to (29) given that in fact (29) is the true regime. Thus (31) can be seen as the probability of a false regime identification. Moreover, in the case that the density of the random variable X_τ admits a closed-form representation, we can also compute

$$\int \mathbb{P}_y^{\tau,(1)}[T(a) = T_D(a)] f_{X_\tau}(y|x) dy = \int \mathbb{P}_y^{\tau,(1)}[T_D(a) \leq T_U(a)] f_{X_\tau}(y|x) dy,$$

which can be seen as the aggregate probability (or unconditional probability) of a false identification for any given change-point τ .




FIG 6. Probability $\mathbb{P}_x[T_D(a) < T_U(b)]$, where $a = b = 1$, as a function of κ/σ^2 , where $\kappa\theta > \sigma^2/2$. Process $\{X_t\}$ is a Cox-Ingersoll-Ross process: $dX_t = \kappa(\theta - X_t)dt + \sigma\sqrt{X_t}dW_t$, $X_0 = x$. We use the same values of parameters as in Figure 4.

6. Conclusion. In this paper, we discussed properties of a diffusion process stopped at time $T_D(a)$, the first time when the process drops by amount a from its running maximum. We derived the joint distribution of the running minimum and the running maximum stopped at time $T_D(a)$. This allowed us to obtain a formula for the probability that a drawdown of size a precedes a drawup of size b .

A possible extension of our work is the calculation of the probability that a drawup precedes a drawdown in a finite time horizon. This would require a combination of our results with the distributions of times $T_D(a)$ and $T_U(b)$. We do not expect this would lead to a closed form solution.

REFERENCES

- [1] BARNARD, G. (1959): Control charts and stochastic processes, *Journal of the Royal Statistical Society: B*, Vol. 21, 239-257.
- [2] BISSELL, A. (1969): Cusum techniques for quality control, *Applied Statistics*, Vol. 18, 1-30.
- [3] CHERNY, A. AND B. DUPIRE (2007): On Certain Distributions Associated with the Range of Martingales, *Preprint*.
- [4] CHEKHOV, A., S. URYASEV, AND M. ZABARANKIN (2005): Drawdown Measure in Portfolio Optimization, *International Journal of Theoretical and Applied Finance*, Vol. 8, No. 1, 13-58.
- [5] COX, J. C., J. E. INGERSOLL, AND S. A. ROSS (1985): A Theory of the Term Structure of Interest Rates, *Econometrica*, Vol. 53, 385-407.
- [6] CVITANIC, J. AND I. KARATZAS (1995): On Portfolio Optimization under Drawdown Constraints, *IMA Lecture Notes in Mathematics & Applications*, Vol. 65, 77-88.
- [7] VAN DOBBEN DE BRUYN, D. S. (1968): Cumulative Sum tests: Theory and Practice, *Hafner, New York*.

- [8] GIHMAN, I. I. AND A. V. SKOROKHOD (1972): Stochastic Differential Equations, Springe-Verlag, New York.
- [9] GROSSMAN, S. J. AND Z. ZHOU (1993): Optimal investment strategies for controlling drawdowns, *Mathematical Finance*, Vol. 3, No. 3, 241-276.
- [10] HADJILIADIS, O. AND G. V. MOUSTAKIDES (2006): Optimal and Asymptotically Optimal CUSUM Rules for Change Point Detection in the Brownian Motion Model with Multiple Alternatives, *Theory of Probability and its Applications*, Vol. 50, No. 1, 131-144.
- [11] HADJILIADIS, O. AND J. VECER (2006): Drawdowns Preceding Rallies in a Brownian Motion Process, *Quantitative Finance*, Vol. 6, No. 5, 403-409.
- [12] KEMP, K. (1961): The average run-length of the cumulative sum chart when a V-mask is used, *Journal of the Royal Statistical Society: B*, Vol. 23, 149-153.
- [13] KHAN, R. A. (2008): Distributional Properties of CUSUM Stopping Times, *Sequential Analysis*, Vol. 27, No. 4, 420-434.
- [14] LEHOCZKY, J. P. (1977): Formulas for Stopped Diffusion Processes with Stopping Times Based on the Maximum, *Annals of Probability*, Vol. 5, No. 4, 601-607.
- [15] MAGDON-ISMAIL, M., A. ATIYA, A. PRATAP, AND Y. ABU-MOSTAFA (2004): On the Maximum Drawdown of a Brownian Motion, *Journal of Applied Probability*, Vol. 41, No. 1.
- [16] MEILIJSON, I. (2003): The Time to a Given Drawdown in Brownian Motion, *Seminaire de Probabilités*, XXXVII, 94-108.
- [17] NIKEGBALI, A. (2006): A Class of Remarkable Submartingales, *Stochastic Processes and their Applications*, Vol. 116, No. 6, 917-938.
- [18] OBLOJ, J. AND M. YOR (2006): On Local Martingale and its Supremum: Harmonic Functions and beyond, *Springer, From Stochastic Calculus to Mathematical Finance*, 517-533.
- [19] POOR, H. V. AND O. HADJILIADIS (2008): Quickest Detection, *Cambridge University Press, Cambridge, UK*.
- [20] SALMINEN, P. AND P. VALLOIS (2007): On Maximum Increase and Decrease of Brownian Motion, *Annales de l'Institut Henri Poincare (B) Probability and Statistics*, Vol. 43, No. 6.
- [21] SHREVE, S. (2004): Stochastic Calculus for Finance II, *Springer Verlag*.
- [22] TAYLOR, H. M. (1975): A Stopped Brownian Motion Formula, *Annals of Probability*, Vol. 3, No. 2, 234-246.
- [23] VASICEK, O. A. (1977): An Equilibrium Characterization of the Term Structure, *Journal of Financial Economics*, Vol. 5, 177-188.
- [24] VECER, J. (2006): Maximum Drawdown and Directional Trading, *Risk*, Vol. 19, No. 12, 88-92.
- [25] WOODALL, W. H. (1984): On the Markov Chain approach to the two-sided CUSUM procedure, *Technometrics*, Vol. 26, 41-46.

DEPARTMENT OF STATISTICS, COLUMBIA UNIVERSITY, 1255 AMSTERDAM AVE, NY 10027, USA,
DEPARTMENT OF MATHEMATICS, BROOKLYN COLLEGE (C.U.N.Y.), INGERSOLL HALL, NY 11209.
E-MAIL: lp2185@columbia.edu; vecer@stat.columbia.edu; ohadjiliadis@brooklyn.cuny.edu